

Research Project

Teturo Kamae

In this research, I'll study the meaning of randomness and quantities representing the degree of randomness. To answer to the philosophical question what is random straightforwardly, the logical point of view is necessary. In fact, randomness is defined using algorithmic notion, Kolmogorov-Chaitin complexity. But, the random numbers thus defined have a fatal weak point, that is, they do exist with probability 1, but none of them are not constructible. On the other hand, pseudo-random numbers are indispensable tool for the computer simulations and they have to be constructed algorithmically. In this research, forgetting the philosophical aspect, we put emphasis on the practical aspect of randomness. I also want to continue another study of the deviation of width of convex sets along random directions.

Normal numbers are defined as infinite sequences realizing the law of large number for the uniform i.i.d. random variables in the sense that the relative frequency of blocks occurring in them coincide with their expectations. In the 1970's, it was asked to what extent the normal numbers are considered to be random from the point view of absence of winning strategy (Von Mises' notion of collective). We choose a subsequence of an infinite sequence looking it up to the $(n - 1)$ -th place to decide whether to put the n -th place into the subsequence or not. If we use a finite automaton for this decision, then it was known that the infinite subsequence obtained from any normal number is again a normal number. Thus, the normality is preserved. I generalized this result for a wider class of infinite automata. I want to revisit this problem to find a necessary and sufficient condition for to preserve the normality.

To be a normal number is a minimum requirement to be a random number, but it is too weak. We need a stronger requirement. For this purpose, I introduced a criterion $\Sigma(x)$ for finite sequences $x = x_1 \cdots x_n \in \mathbb{A}^n$ over an alphabet \mathbb{A} ($2 \leq d := \#\mathbb{A} < \infty$) to measure the degree of randomness. It is defined as the sum over all finite blocks ξ so that $\Sigma(x) = \sum_{\xi} |x_1 \cdots x_n|_{\xi}^2$, where $|x_1 \cdots x_n|_{\xi}$ is the number of oc-

currences of ξ in x . Comparing this value among finite sequences of the same length, smaller values imply more random. It is proved that $\lim_{n \rightarrow \infty} (1/n^2) \Sigma(X_1 \cdots X_n) = \frac{d+1}{2(d-1)}$ holds with probability 1 for the uniform i.i.d. random variables X_1, X_2, \dots over \mathbb{A} . I call the infinite sequences $x_1 x_2 \cdots$ satisfying this almost all property Σ -random numbers. This notion is stronger than the normal numbers. We know an algorithm to construct a Σ -random number starting from any finite sequences. I want to study the properties of the Σ -random numbers to see how they are good as pseudo-random numbers. Also, for a 2-dimensional configuration x , we want to discuss the randomness by a 2-dimensional version of $\Sigma(x)$.

I also want to continue a joint work with Prof. Akiyama (Tsukuba University) to study the deviation of the width of convex sets in the plain or the 3-dimensional space along random directions. We proved that among the n -gons with odd n in the plain the deviation attains minimum if and only if it is the regular n -gon. If n is even, we obtained the n -gon attaining the minimum if n is not a power of 2, which is not the regular one. It is still open in the case of a power of 2. In this research, we try to solve this open problem as well as developing the study to the 3-dimensional space.

Recently I started studies of the stream zeros Ω_P determined by a primitive polynomial $P(z) = a_k z^k + \cdots + a_1 z + a_0$ of integral coefficients, that is

$$\Omega_P = \{(x_n \in R/Z; n \in Z); a_k x_{n-k} + \cdots + a_1 x_{n-1} + a_0 x_n = 0 (\forall n \in Z)\},$$

from the point views of dynamical systems and algebra. The results obtained at this moment are on the paper [81] in the list of paper. I want to develop this research.