

Free ribbon lemma for surface-link

Akio Kawauchi

Osaka Central Advanced Mathematical Institute, Osaka Metropolitan University

Sugimoto, Sumiyoshi-ku, Osaka 558-8585, Japan

kawauchi@omu.ac.jp

ABSTRACT

A free surface-link is a surface-link whose fundamental group is a free group not necessarily meridian-based. Free ribbon lemma says that every free sphere-link in the 4-sphere is a ribbon sphere-link. Four different proofs of Free ribbon lemma are explained. The first proof is done in an earlier paper. The second proof is done by showing that there is an O2-handle basis of a ribbon surface-link. The third proof is done by removing the commuter relations from a Wirtinger presentation of a free group, which a paper on another proof of Free ribbon lemma complements. The fourth proof is given by the special case of the proof of the result that every free surface-link is a ribbon surface-link which is a stabilization of a free ribbon sphere-link. As a consequence, it is shown that a surface-link is a sublink of a free surface-link if and only if it is a stabilization of a ribbon sphere-link.

Keywords: Free ribbon lemma, Free surface-link, Ribbon sphere-link, Stabilization.

Mathematics Subject Classification 2010: Primary 57N13; Secondary 57Q45

1. Introduction

A *surface link* is a closed, possibly disconnected, oriented surface F smoothly embedded in the 4-sphere S^4 , and it is called a *surface knot* if F is connected. If F consists of 2-spheres F_i ($i = 1, 2, \dots, r$), then F is called a *sphere-link* (or an S^2 -link) of r components. It is shown that a surface-link F is a trivial surface-link (i.e., bounds disjoint handlebodies in S^4) if the fundamental group $\pi_1(S^4 \setminus F, x_0)$ is a meridian-based free group, [10], [11], [12]. A surface-link F is *ribbon* if F is obtained from

a trivial S^2 -link O in S^4 by surgery along a smoothly embedded disjoint 1-handle system h^O on O , [20], [29], [30], [32]. A surface-link F in the 4-sphere S^4 is *free* if the fundamental group $\pi_1(S^4 \setminus F, x_0)$ is a (not necessarily meridian-based) free group. In this paper, four different proofs of the following *Free ribbon lemma* and its generalization to a general free surface-link are explained.

Free ribbon lemma. Every free S^2 -link in S^4 is a ribbon S^2 -link.

Free ribbon lemma leads to the following conjectures: Poincaré conjecture, [15], [24], [25], [26]. J. H. C. Whitehead asphericity conjecture for aspherical 2-complex, [5], [14], [17], [28]. Kervaire conjecture on group weight, [1], [16], [21], [23], [22]. The first proof is given [14]. For convenience, an outline of the first proof is explained here.

First proof of Free ribbon lemma. Let L_i ($i = 1, 2, \dots, r$) be the components of a free S^2 -link L in S^4 . By a base change of the free fundamental group $\pi_1(S^4 \setminus L, x_0)$, take a basis x_i ($i = 1, 2, \dots, r$) of $\pi_1(S^4 \setminus L, x_0)$ inducing a meridian basis of L in $H_1(S^4 \setminus L; \mathbb{Z})$, [23]. Let Y be the 4-manifold obtained from S^4 by surgery along L , which is diffeomorphic to the connected sum of r copies $S^1 \times S_i^3$ ($i = 1, 2, \dots, r$) of $S^1 \times S^3$, [13], [14]. Under a canonical isomorphism $\pi_1(S^4 \setminus L, x_0) \rightarrow \pi_1(Y, x_0)$, the factors $S^1 \times p_i$ ($i = 1, 2, \dots, r$) of $S^1 \times S_i^3$ ($i = 1, 2, \dots, r$) with suitable paths to the base point x_0 represent the basis x_i ($i = 1, 2, \dots, r$). Let k_i ($i = 1, 2, \dots, r$) be the loop system in Y produced from the components L_i ($i = 1, 2, \dots, r$) by the surgery. By using the fact that any homotopy deformations of k_i ($i = 1, 2, \dots, r$) in Y do not change the link type of the surface-link L in S^4 , the loop system k_i ($i = 1, 2, \dots, r$) is homotopically deformed in Y so that the surface-link L in S^4 obtained from the deformed loop system k_i ($i = 1, 2, \dots, r$) by back surgery is a ribbon surface-link in S^4 , completing the proof of Free ribbon lemma.

To explain the second and third proofs of Free ribbon lemma, the notion of an O2-handle basis of a surface-link is needed, [10], [19]. An *O2-handle pair* on a surface-link F in S^4 is a pair $(D \times I, D' \times I)$ of 2-handles $D \times I$, $D' \times I$ on F in S^4 which intersect orthogonally only with the attaching parts $(\partial D) \times I$, $(\partial D') \times I$ to F , so that the intersection $Q = (\partial D) \times I \cap (\partial D') \times I$ is a square. Let $(D \times I, D' \times I)$ be an O2-handle pair on a surface-link F . Let $F(D \times I)$ and $F(D' \times I)$ be the surface-links obtained from F by the surgeries along $D \times I$ and $D' \times I$, respectively. Let $F(D \times I, D' \times I)$ be the surface-link which is the union $\delta \cup F_\delta^c$ of the plumbed disk

$$\delta = \delta_{D \times I, D' \times I} = D \times \partial I \cup Q \cup D' \times \partial I$$

and the surface $F_\delta^c = \text{cl}(F \setminus (\partial D \times I \cup \partial D' \times I))$. The surface-links $F(D \times I)$, $F(D' \times I)$ and $F(D \times I, D' \times I)$ are equivalent surface-links, [10]. An *O2-handle basis* of a surface-link F is a disjoint system of O2-handle pairs $(D_i \times I, D'_i \times I)$ ($i = 1, 2, \dots, r$) on F in S^4 such that the boundary loop pair system $(\partial D_i, \partial D'_i)$ ($i = 1, 2, \dots, r$) of the core disk system (D_i, D'_i) ($i = 1, 2, \dots, r$) of $(D_i \times I, D'_i \times I)$ ($i = 1, 2, \dots, r$) is a spin loop basis for F in S^4 , which is a system of a spin loop basis of every component F_i of F . Note that there is a spin loop basis for every surface-knot in F , [3]. In this paper, for simplicity, an O2-handle basis $(D_i \times I, D'_i \times I)$ ($i = 1, 2, \dots, r$) for F is denoted by $(D \times I, D' \times I)$. The surgery surface-link of F by $(D_i \times I, D'_i \times I)$ ($i = 1, 2, \dots, r$) is denoted by $F(D \times I, D' \times I)$. The following theorem is shown for the second and third proofs of Free ribbon lemma.

Theorem 1.1. For every free ribbon surface-link F in S^4 , there is an O2-handle basis $(D \times I, D' \times I)$ on F in S^4 such that $D \times I$ belongs to the 1-handle system of the ribbon surface-link F .

The second proof of Free ribbon lemma is explained as follows.

Second proof of Free ribbon lemma. Let L be a free S^2 -link. Then there is a ribbon surface-link F such that the fundamental group $\pi_1(S^4 \setminus F, x_0)$ is isomorphic to the free fundamental group $\pi_1(S^4 \setminus L, x_0)$ by a meridian-preserving isomorphism, [18]. By Theorem 1.1, the surgery surface-link $L' = F(D \times I, D' \times I)$ is a ribbon S^2 -link, [10], [19]. Then there is a meridian-preserving isomorphism $\pi_1(S^4 \setminus L', x_0) \rightarrow \pi_1(S^4 \setminus L, x_0)$ on free groups, which implies that L' is equivalent to L , [14], [18]. Thus, L is a ribbon S^2 -link, completing the proof of Free ribbon lemma.

The third proof of Free ribbon lemma is related to a Wirtinger presentation of a free group. A finite group presentation $(x_1, x_2, \dots, x_n \mid R_1, R_2, \dots, R_m)$ is a Wirtinger presentation if $R_j = W_j x_{s_j} W_j^{-1} x_{t_j}^{-1}$ for some indexes s_j, t_j in $\{1, 2, \dots, n\}$ for every j ($j = 1, 2, \dots, m$). The relator R_j is a *commutator relation* if $x_{s_j} = x_{t_j}$. It is well-known that a Wirtinger presentation of a finitely presented group G with $H_1(G; \mathbb{Z}) \cong \mathbb{Z}^r$ is always equivalent (without changing the generating set) to a Wirtinger presentation P such that the Wirtinger presentation P' obtained by removing all the commutator relations from P has deficiency r . Such a Wirtinger presentation P is called a *normal* Wirtinger presentation. The following corollary is obtained from Theorem 1.1.

Corollary 1.2. If a free group G of rank r has a normal Wirtinger presentation P , then G has the Wirtinger presentation P' of deficiency r obtained from P by removing

all the commutator relations.

Proof of Corollary 1.2 assuming Theorem 1.1. For a free group G of rank r , let $P = (x_1, x_2, \dots, x_n | R_1, R_2, \dots, R_m)$ be a normal Wirtinger presentation of G such that the relators R_j ($n - r + 1 \leq j \leq m$) are the commutator relations. Let O be a trivial S^2 -link of n components in S^4 such that the meridian basis of the free fundamental group $\pi_1(S^4 \setminus O, x_0)$ are identified with x_i ($i = 1, 2, \dots, n$). Let h_j ($1 \leq j \leq m$) be the 1-handles on O indicated by the relators R_j ($1 \leq j \leq m$). By the van Kampen theorem, the ribbon surface-link F in S^4 obtained by surgery along h_j ($1 \leq j \leq m$) has the normal Wirtinger presentation P of the fundamental group $\pi_1(S^4 \setminus F, x_0)$ with the meridian generators set $\{x_1, x_2, \dots, x_n\}$, [7], [8]. Let L be the ribbon surface-link obtained from O by surgery along the 1-handles h_j ($1 \leq j \leq n - r$), which is a ribbon S^2 -link of r components. The fundamental group $\pi_1(S^4 \setminus L, x_0)$ has the Wirtinger presentation P' of deficiency r obtained from P by removing all the commutator relations. By Theorem 1.1, the 1-handles h_j ($n - r + 1 \leq j \leq m$) on L are trivial 1-handles, so that $\pi_1(S^4 \setminus L, x_0)$ is isomorphic to $\pi_1(S^4 \setminus F, x_0)$ by a meridian-preserving isomorphism. This completes the proof of Corollary 1.2 assuming Theorem 1.1.

The author has published a paper on another proof of Free ribbon lemma, which this paper complements, [18]. The third proof of Free ribbon lemma is nothing but the proof of the paper except for adding to it the assertion of Corollary 1.5 which was missing from it. For convenience, an outline of the third proof is explained here.

Third proof of Free ribbon lemma. Let L be a free S^2 -link of r components. Since the fundamental group $G = \pi_1(S^4 \setminus L, x_0)$ is a free group with $H_1(G; \mathbb{Z}) = \mathbb{Z}^r$ and $H_2(G; \mathbb{Z}) = 0$, there is a normal Wirtinger presentation P of G whose generator set comes from meridians of L in S^4 , [18], [31]. Note that there is also another method to find such a normal Wirtinger presentation P using a normal form of L in S^4 , [6], [7], [8], [20]. Let L' be a ribbon S^2 -link given by the Wirtinger presentation P' obtained from P by removing all the commutators. By Corollary 1.2, there is a meridian-preserving isomorphism $\pi_1(S^4 \setminus L', x_0) \rightarrow \pi_1(S^4 \setminus L, x_0)$, so that L' is equivalent to L . Thus, L is a ribbon S^2 -link, completing the proof of Free ribbon lemma.

The fourth proof of Free ribbon lemma is given by a direct proof of the following theorem.

Theorem 1.3. Every free surface-link F in S^4 is a ribbon surface-link in S^4 .

Fourth proof of Free ribbon lemma. It is obtained from Theorem 1.3 by restricting F to every free S^2 -link, completing the proof of Free ribbon lemma.

Thus, after the proofs of Theorems 1.1 and 1.3, there are four different proofs of Free ribbon lemma. To generalize the free ribbon lemma to a free surface-link, the notion of a stabilization of a surface-link is needed, [10], [19]. A *stabilization* of a surface-link L is a connected sum $F = L \#_{k=1}^s T_k$ of L and a system of trivial torus-knots T_k ($k = 1, 2, \dots, s$). By granting $s = 0$, a surface-link L itself is regarded as a stabilization of L . Free ribbon lemma is generalized to a general free surface-link as follows.

Corollary 1.4. Every free surface-link F in S^4 is a stabilization of a free ribbon S^2 -link L in S^4 .

Proof of Corollary 1.4 assuming Theorems 1.1 and 1.3. Theorems 1.1 and 1.3 imply that every free surface-link F is a ribbon surface-link and a stabilization of a free S^2 -link L , [10]. By Free ribbon lemma, the free S^2 -link L is a ribbon S^2 -link. This result also follows directly, [19]. This completes the proof of Corollary 1.4 assuming Theorems 1.1 and 1.3.

It is shown that an S^2 -link L is a sublink of a free S^2 -link if and only if L is a ribbon S^2 -link, [14]. The following corollary generalizes this property to a general surface-link.

Corollary 1.5. A surface-link L in S^4 is a sublink of a free surface-link F in S^4 if and only if L is a stabilization of a ribbon S^2 -link in S^4 .

Proof of Corollary 1.5 assuming Corollary 1.4. If L is a sublink of a free surface-link F , then L is a stabilization of a ribbon S^2 -link since every free surface-link is a stabilization of a free ribbon S^2 -link by Corollary 1.4. Conversely, if L is a stabilization of a ribbon S^2 -link, then L is a sublink of a stabilization of a free ribbon S^2 -link which is a free surface-link F since every ribbon S^2 -link is a sublink of a free S^2 -link. This completes the proof of Corollary 1.5 assuming Corollary 1.4.

2. Proofs of Theorems 1.1 and 1.3.

Let F be a *free* surface-link in S^4 with components F_i ($i = 1, 2, \dots, r$). Let $N(F) = \cup_{i=1}^r N(F_i)$ be a tubular neighborhood of $F = \cup_{i=1}^r F_i$ in S^4 which is a trivial normal disk bundle $F \times D^2$ over F , where D^2 denotes the unit disk of complex numbers of norm ≤ 1 . Let $E = E(F) = \text{cl}(S^4 \setminus N(F))$ be the exterior of F in S^4 . The boundary

$\partial E = \partial N(F) = \cup_{i=1}^r \partial N(F_i)$ of the exterior E is a trivial normal circle bundle over $F = \cup_{i=1}^r F_i$. Identify $\partial N(F_i) = F_i \times S^1$ for $S^1 = \partial D^2$ such that the composite inclusion

$$F_i \times 1 \rightarrow \partial N(F_i) \rightarrow \text{cl}(S^4 \setminus N(F_i))$$

induces the zero-map in the integral first homology. The following lemma uses the assumption that the fundamental group $\pi_1(E, x_0)$ is a free group of rank r and the fact that the first homology group $H_1(E; Z)$ is a free abelian group of rank r with meridian basis.

Lemma 2.1. The composite inclusion $F_i \times 1 \rightarrow \partial N(F_i) \rightarrow E$ is null-homotopic for all i .

Proof of Lemma 2.1. Since $\partial N(F_i) = F_i \times S^1$, the fundamental group elements between the factors $F_i \times 1$ and $q_i \times S^1$ are commutative. Let $a_i (i = 1, 2, \dots, r)$ be embedded edges with common vertex x_0 in E such that $a_i \setminus \{x_0\} (i = 1, 2, \dots, r)$ are mutually disjoint and $a_i \cap (\cup_{j=1}^r F_j \times 1) = p_i \times 1$ for a point p_i of $F_i \times 1$. The surface $F_i \times 1$ in $\partial N(F_i) = F_i \times S^1$ is chosen so that the inclusion $F_i \times 1 \rightarrow \text{cl}(S^4 \setminus N(F_i))$ induces the zero-map in the integral first homology. Since $H_1(E; Z)$ is a free abelian group of rank r with meridian basis and $\pi_1(E, x_0)$ is a free group of rank r , the image of the homomorphism $\pi_1(a_i \cup F_i \times S^1, x_0) \rightarrow \pi_1(E, x_0)$ is an infinite cyclic group generated by the homotopy class $[a_i \cup p_i \times S^1]$. This implies that the inclusion $F_i \times 1 \rightarrow E$ is null-homotopic. This completes the proof of Lemma 2.1. \square

By using the free group $\pi_1(E, x_0)$ of rank r , let $\Gamma = \cup_{i=1}^r a_i \cup C_i$ be a connected graph in the interior $\text{Int}(E)$ of E consisting of embedded edges $a_i (i = 1, 2, \dots, r)$ with the common base point x_0 and disjoint embedded circles $C_i (i = 1, 2, \dots, r)$ such that

- (1) the half-open edges $a_i \setminus \{x_0\} (i = 1, 2, \dots, r)$ are mutually disjoint and $a_i \cap (\cup_{j=1}^r C_j) = v_i$, a point in C_i for every i ,
- (2) the inclusion $i : (\Gamma, x_0) \rightarrow (E, x_0)$ induces an isomorphism $i_{\#} : \pi_1(\Gamma, x_0) \rightarrow \pi_1(E, x_0)$, and
- (3) the homology class $[p_i \times S^1] = [C_i]$ in $H_1(E; Z)$ for all i .

In fact, by (2), the homotopy classes $[a_i \cup C_i] (i = 1, 2, \dots, r)$ form a basis of the free group $\pi_1(E, q_0)$. (3) is obtained by a base change of the free group $\pi_1(E, x_0)$, [23]. Since Γ is a $K(\pi, 1)$ -space, there is a piecewise-linear map $f : (E, x_0) \rightarrow (\Gamma, x_0)$ inducing the inverse isomorphism $f_{\#} = (i_{\#})^{-1} : \pi_1(E, q_0) \rightarrow \pi_1(\Gamma, q_0)$ of $i_{\#}$, and by

the homotopy extension property, the restriction of f to Γ is the identity map, [27]. The restriction of f to ∂E is homotopic to the composite map

$$g : \partial E = F \times S^1 \rightarrow \cup_{i=1}^r q_i \times S^1 \rightarrow \Gamma$$

such that the first map $F \times S^1 \rightarrow \cup_{i=1}^r q_i \times S^1$ is induced from the constant map $F \rightarrow \cup_{i=1}^r \{q_i\}$ and the second map $\cup_{i=1}^r q_i \times S^1 \rightarrow \Gamma$ is defined by the map f . By using a boundary collar of ∂E in E , assume that the piecewise-linear map $f : (E, x_0) \rightarrow (\Gamma, x_0)$ defines the map $g : \partial E \rightarrow \Gamma$. For a non-vertex point p_i of C_i , the preimage $V_i = (f)^{-1}(p_i)$ is a bi-collard compact oriented proper piecewise-linear 3-manifold in E . Take the compact 4-manifold E' obtained from E by splitting along $V = \cup_{i=1}^r V_i$ to be connected. Then join the components in each V_i with 1-handles in E' . By these modifications, V_i is assumed to be connected for all i within a homotopic deformation of f . The boundary ∂V_i is the disjoint union $P_i(F)$ of m_{ij} parallel copies $m_{ij}F_j$ of $F_j \times 1$ for all j ($j = 1, 2, \dots, r$) in S^4 , where m_{ii} is an odd integer and m_{ij} for distinct i, j is an even integer. Let $P(F) = \cup_{i=1}^r P_i(F)$ be the surface-link in S^4 . Let h_i be a disjoint 1-handle system on $P_i(F)$ embedded in V_i such that the surface $P_i(F; h_i)$ obtained from $P_i(F)$ by surgery along h_i is connected and the genus of $P_i(F; h_i)$ is equal to the total genus of $P_i(F)$. Assume that one copy of the parallel $m_{ii}F_i$ of F_i is identified with F_i and just one 1-handle h_i^F of h_i attaches to F_i . Let $P(F; h) = \cup_{i=1}^r P_i(F; h_i)$ be a surface-link in S^4 . By further taking a disjoint 1-handle system h'_i on $P_i(F; h_i)$ embedded in V_i , the closed surface $P_i(F; h_i, h'_i)$ obtained from $P_i(F; h_i)$ by surgery along h'_i bounds a handlebody V'_i in V_i , so that the surface-link $P(F; h, h') = \cup_{i=1}^r P_i(F; h_i, h'_i)$ is a trivial surface-link in S^4 . Because of the isomorphism $f_\#$, the compact 4-manifold E' is simply connected, so that the 1-handle system $h' = \cup_{i=1}^r h'_i$ is a trivial 1-handle system on the surface-link $P(F; h)$ in S^4 , [4], [12]. Thus, the surface-link $P(F; h)$ is a trivial surface-link in S^4 , [10], [11]. The proof of Theorem 1.1 is done as follows.

Proof of Theorem 1.1. A ribbon surface-link F is obtained from a trivial S^2 -link O in S^4 by surgery along a disjoint 1-handle system h^O on O , so that the surface-link $P(F)$ of a free ribbon surface-link F is a ribbon surface-link obtained from a trivial S^2 -link $P(O)$ in S^4 by surgery along a disjoint 1-handle system $P(h^O)$ on $P(O)$. Let $VP(F)$ be a multi-punctured handlebody system with $\partial VP(F) = P(F) \cup P(O)$ constructed from the collar $P(O) \times [0, 1]$ of O in S^4 by attaching the 1-handle system $P(h^O)$ to $P(O) \times 0 = P(O)$ which is called a SUPH system for the ribbon surface-link $P(F)$ in S^4 , [19]. The 1-handle system $h = \cup_{i=1}^r h_i$ on $P(F)$ and the SUPH system $VP(F)$ construct a SUPH system $VP(F) \cup h$ for the trivial surface-link $P(F; h)$ with $\partial(VP(F) \cup h) = P(F; h) \cup P(O)$. A spin loop basis (ℓ, ℓ') for $P(F)$ is the system consisting of a spin loop basis of every component of $P(F)$ where the spin

loop system ℓ belongs to a meridian system of the 1-handle system $P(h^O)$. This loop system (ℓ, ℓ') is a spin loop basis of the trivial ribbon surface-link $P(F; h)$. Equivalent ribbon surface-links are faithfully equivalent, and they are moved into each other by the moves M_0, M_1, M_2 , [9]. This means that there is an orientation-preserving diffeomorphism w of S^4 sending a multi-punctured manifold of the SUPH system $VP(F) \cup h$ for $P(F; h)$ to a multi-punctured manifold of the standard handlebody system W in S^4 . By a choice of w , the system $(w(\ell), w(\ell'))$ is a meridian-longitude pair system of W in S^4 , [2], [10]. The loop system $w(\ell')$ bounds a disjoint disk system δ' in S^4 with $\delta' \cap W = w(\ell')$, so that the loop system ℓ' bounds a disjoint disk system $D' = w^{-1}(\delta')$ in S^4 with $D' \cap (VP(F) \cup h) = \ell'$. The loop system ℓ belongs to a meridian system of the 1-handle system $P(h^O)$ and hence bounds a subsystem D of the meridian disk system $P(h^O)$. Thus, it is shown that there is an O2-handle basis $(D \times I, D' \times I)$ on $P(F)$ in S^4 , whose subsystem to F gives an O2-handle basis on F in S^4 . This completes the proof of Theorem 1.1.

The proof of Theorem 1.3 is done as follows.

Proof of Theorem 1.3. An *anti-parallel surface-link* of F_j in S^4 is the boundary surface-link $\partial(F_j \times J)$ of a normal line bundle $F_j \times J$ of F_j in S^4 such that the natural homomorphism $H_1(F \times 1; Z) \rightarrow H_1(S^4 \setminus F \times 0; Z)$ is the zero-map, where $J = [0, 1]$. Note that the boundary surface-knot $\partial(F_j^{(0)} \times J)$ for a compact once-punctured surface $F_j^{(0)}$ of F_j is a trivial surface-knot in S^4 since $F_j^{(0)} \times J$ is a handlebody. Let $n_{ii} = (m_{ii} - 1)/2$ and $n_{ij} = m_{ij}/2$ for distinct i, j . The surface-link $P_i(F)$ in V_i consists of F_i and anti-parallel surface-links of F_j in $P_i(F)$ for all j , denoted by $p_{ijk}(F)$ ($k = 1, 2, \dots, n_{ij}; j = 1, 2, \dots, r$). Since E' is simply connected, the 1-handle system h_i can be chosen so that a trivial surface-knot F_{ijk} is obtained from the anti-parallel surface-link $p_{ijk}(F)$ in $P_i(F)$ by surgery along a unique 1-handle h_{ijk} in h_i . Let $h(0)$ be the system of the 1-handles h_{ijk} ($k = 1, 2, \dots, n_{ij}; i, j = 1, 2, \dots, r$) in h , and $h(1)$ the complementary system of $h(0)$ in h . Let $P(F; h(0))$ be the surface-link obtained from $P(F)$ by surgery along $h(0)$, which consists of the surface-link F and the trivial surface-knots F_{ijk} ($k = 1, 2, \dots, n_{ij}; i, j = 1, 2, \dots, r$). The trivial surface-link $P(F; h)$ is obtained from the surface-link $P(F; h(0))$ by surgery along the 1-handle system $h(1)$. The surface-link $P(F; h, h')$ bounds a handlebody system V' in V . Note that h' is a trivial 1-handle system on $P(F; h)$. Then there is a disjoint handlebody system U in S^4 with $\partial U = P(F; h)$ extending the handlebody system V' by adding a 2-handle system $e \times I$ which makes an O2-handle system together with a thickened meridian disk system $m(h')$ of h' , [10], [11]. Let d be a meridian disk system of the 1-handle system $h(1)$. In general, the disk system d meets the core disk system e of $e \times I$ transversely in finite points in S^4 , but the handlebody system U is isotopically

deformed so to have $d \cap U = \partial d$ by an isotopic deformation of U in a neighborhood of the 3-disk system $e \times I \cup m(h') \times I$ in S^4 . For $h(1) = d \times I$, the union $U \cup h(1)$ is a compact oriented 3-manifold with boundary $P(F; h(0))$ obtained from U by adding the 2-handle system $h(1)$ since $h(1) \cap U = h(1) \cap \partial U = (\partial d) \times I$. Let (ℓ, ℓ') be a spin loop basis of $P(F; h)$ given for $P(F; h(0))$ such that when restricted to every trivial surface-knot F_{ijk} it becomes a standard spin loop basis. Since $P(F; h)$ is a trivial surface-link, there is an orientation-preserving diffeomorphism w of S^4 sending $P(F; h)$ to the boundary ∂W of a standard handlebody system W in S^4 such that the spin loop basis $(w(\ell), w(\ell'))$ of ∂W is a meridian-longitude pair system of W , [2], [10]. Let $U(W) = w^{-1}(W)$ be the handlebody system in S^4 with $\partial U(W) = P(F; h)$. The spin loop basis $(w(\ell), w(\ell'))$ of ∂W bounds a core disk-pair system (δ, δ') of an O2-handle basis $(\delta \times I, \delta' \times I)$ of the trivial surface-link ∂W in S^4 , where δ denotes a meridian disk system of W . Thus, the spin loop basis (ℓ, ℓ') of $P(F; h)$ bounds the core disk pair system (D, D') of the O2-handle basis $(D \times I, D' \times I) = (w^{-1}(\delta) \times I, w^{-1}(\delta') \times I)$ on $P(F; h)$ in S^4 with D in $U(W)$, so that $d \cap D = h(1) \cap D = \emptyset$. The handlebody system $U(W)$ is isotopically deformed to be $U(W) = U$ in S^4 . Consider the 3-manifold $U' \cup h(1)$ obtained from the 3-manifold $U \cup h(1)$ by splitting along the disk system D , which is a multi-punctured 3-sphere system. Since $\partial(U \cup h(1)) = P(F; h(0))$, the surgery of $P(F; h(0))$ along the 2-handle system $D \times I$ gives that the boundary $\partial(U' \cup h(1))$ consists of the S^2 -link S obtained from F and the S^2 -knots S_{ijk} ($k = 1, 2, \dots, n_{ij}$) for all i, j obtained from the trivial surface-knots F_{ijk} ($k = 1, 2, \dots, n_{ij}$) for all i, j . The S^2 -knots S_{ijk} for all i, j, k are shown to form a trivial S^2 -link in S^4 . To see this, consider a collar $V \times J$ of V in S^4 to move (via a boundary collar of V) the anti-parallel surface $p_{ijk}(F)$ and the 1-handle h_{ijk} into a disjoint subcollar $V \times J_{ijk}$ of $V \times J$ for each i, j, k . Then by forgetting the 1-handle system $h(1)$, the trivial surface-knots F_{ijk} for all i, j, k except for F in the boundary of $U \cup h(1)$ are isotopic to the boundaries of disjoint handlebodies U_{ijk} for all i, j, k in S^4 . For the meridian disk system d of the 1-handle system $h(1)$ on $P(F; h(0))$, there is a proper disk system d_U in U with $\partial d_U = \partial d$ which splits U into disjoint handlebodies U^0 and U_{ijk}^0 for all i, j, k where the boundaries ∂U^0 and ∂U_{ijk}^0 contain the multi-punctured surfaces of F and F_{ijk} , respectively and U_{ijk}^0 contains the subsystem D_{ijk} for F_{ijk} of the 2-handle core disk system D on $P(F; h(0))$. Deform the handlebody U_{ijk}^0 into the handlebody U_{ijk} for all i, j, k , so that the S^2 -knots S_{ijk} for all i, j, k are isotopic to the boundaries of the disjoint 3-balls U'_{ijk} for all i, j, k obtained from the disjoint handlebodies U_{ijk} for all i, j, k by splitting along the disk systems D_{ijk} for all i, j, k . Thus, the system of the S^2 -knots S_{ijk} for all i, j, k forms a trivial S^2 -link in S^4 , as desired. The multi-punctured 3-sphere system $U' \cup h(1)$ then means that the S^2 -link S is a ribbon S^2 -link in S^4 , [14]. This means that the surface-link F is a ribbon surface-link in S^4 because F is obtained from the ribbon S^2 -link S by surgery along

a 1-handle system. This completes the proof of Theorem 1.3.

Acknowledgements. The author has tried to find a simple algebraic method to prove Corollary 1.2 but failed. He is grateful to Igor Mineyev for asking questions to this algebraic method. This work was partly supported by JSPS KAKENHI Grant Number JP21H00978 and MEXT Promotion of Distinctive Joint Research Center Program JPMXP0723833165 and Osaka Metropolitan University Strategic Research Promotion Project (Development of International Research Hubs).

References

- [1] F. González Acuña and A. Ramírez, A knot-theoretic equivalent of the Kervaire conjecture, *J. Knot Theory Ramifications* 15 (2006), 471-478.
- [2] S. Hirose, On diffeomorphisms over surfaces trivially embedded in the 4-sphere, *Algebraic and Geometric Topology* 2 (2002), 791-824.
- [3] J. A. Hillman and A. Kawauchi, Unknotting orientable surfaces in the 4-sphere, *J. Knot Theory Ramifications* 4 (1995), 213-224.
- [4] F. Hosokawa and A. Kawauchi, Proposals for unknotted surfaces in four-space, *Osaka J. Math.* 16 (1979), 233-248.
- [5] J. Howie, Some remarks on a problem of J. H. C. Whitehead, *Topology* 22 (1983), 475-485.
- [6] S. Kamada, Wirtinger presentations for higher dimensional manifold knots obtained from diagrams, *Fund. Math.* 168(2001), 105-112.
- [7] A. Kawauchi, On the surface-link groups, *Intelligence of low dimensional topology 2006*, Series on knots and everything 40 (2007), 157-164, World Sci. publ.
- [8] A. Kawauchi, The first Alexander $\mathbb{Z}[\mathbb{Z}]$ -modules of surface-links and of virtual links, *Geometry & Topology Monographs* 14 (2008), 353-371.
- [9] A. Kawauchi, Faithful equivalence of equivalent ribbon surface-links, *Journal of Knot Theory and Its Ramifications* 27 (2018), 1843003 (23 pages).
- [10] A. Kawauchi, Ribbonness of a stable-ribbon surface-link, I. A stably trivial surface-link, *Topology and its Applications* 301 (2021), 107522 (16pages).

- [11] A. Kawauchi, Uniqueness of an orthogonal 2-handle pair on a surface-link, *Contemporary Mathematics (UWP)* 4 (2023), 182-188.
- [12] A. Kawauchi, Triviality of a surface-link with meridian-based free fundamental group, *Transnational Journal of Mathematical Analysis and Applications* 11 (2023), 19-27.
- [13] A. Kawauchi, Smooth homotopy 4-sphere, *WSEAS Transactions on Mathematics* 22 (2023), 690-701.
- [14] A. Kawauchi, Ribbonness of Kervaire's sphere-link in homotopy 4-sphere and its consequences to 2-complexes, *J Math Techniques Comput Math* 3(4) (2024), 01-08 (online).
- [15] A. Kawauchi, Classical Poincaré conjecture via 4D topology, *J Math Techniques Comput Math* 3(4) (2024), 1-7 (online).
- [16] A. Kawauchi, Kervaire conjecture on weight of group via fundamental group of ribbon sphere-link, *J Math Techniques Comput Math* 3(4) (2024), 1-3 (online).
- [17] A. Kawauchi, Whitehead aspherical conjecture via ribbon sphere-link, *J Math Techniques Comput Math* 3(5) (2024), 01-10 (online).
- [18] A. Kawauchi, Another proof of free ribbon lemma, *J Math Techniques Comput Math* 3(9) (2024), 01-03 (online).
- [19] A. Kawauchi, Ribbonness of a stable-ribbon surface-link, II. General case (MDPI) *Mathematics* 13 (3), 402 (2025), 1-11. <https://doi.org/10.3390/math13030402>.
- [20] A. Kawauchi, T. Shibuya, S. Suzuki, Descriptions on surfaces in four-space I : Normal forms, *Mathematics Seminar Notes, Kobe University* 10 (1982), 75-125; II: Singularities and cross-sectional links, *Mathematics Seminar Notes, Kobe University* 11 (1983), 31-69. <https://sites.google.com/view/kawauchiwriting>.
- [21] M. A. Kervaire, On higher dimensional knots, *Differential and combinatorial topology* 27 (1965), 105-119, Princeton Univ. Press.
- [22] Ant. A. Klyachko, A funny property of a sphere and equations over groups, *Comm. Algebra* 21 (1993), 2555-2575.
- [23] W. Magnus, A. Karrass and D. Solitar, *Combinatorial group theory: Presentations of groups in terms of generators and relations*, Interscience Publishers (1966).

- [24] G. Perelman, Ricci flow with surgery on three-manifolds. arXiv: math.DG/0303109 v1, 10 Mar 2003.
- [25] H. Poincaré, Second complément à l'Analysis Situs, Proc. London Math. Soc. 32 (1900), 277-308.
- [26] H. Poincaré, Cinquième complément à l'Analysis Situs, Rend. Circ. Mat. Palermo 18 (1904), 45-110.
- [27] E. H. Spanier, Algebraic topology, MacGraw Hill (1966).
- [28] J. H. C. Whitehead, On adding relations to homotopy groups, Ann. Math. 42 (1941), 409-428.
- [29] T. Yajima, On the fundamental groups of knotted 2-manifolds in the 4-space, J. Math. Osaka City Univ. 13 (1962), 63-71.
- [30] T. Yajima, On simply knotted spheres in R^4 , Osaka J. Math. 1 (1964), 133-152.
- [31] T. Yajima, Wirtinger presentations of knot groups, Proc. Japan Acad. 46 (1970), 997-1000.
- [32] T. Yanagawa, On ribbon 2-knots; the 3-manifold bounded by the 2-knot, Osaka J. Math. 6 (1969), 447-164.