

Plan of the study (Yosuke Saito)

For a complex number q satisfying $|q|<1$ and $x\in\mathbb{C}$, set $(x; q)_\infty:=\prod_{n\geq 0}(1-xq^n)$ (q -infinite product). For $x\in\mathbb{C}\setminus\{0\}$, set $\Theta_q(x):=(q; q)_\infty(x; q)_\infty(qx^{-1}; q)_\infty$ (theta function). By setting $D_x=x\frac{\partial}{\partial x}$ (Euler derivative), we define $E_k(x; q):=-D_x^k\log\Theta_q(x)$ ($k\in\mathbb{Z}_{>0}$). For complex numbers q, p satisfying $|q|<1, |p|<1$ and $x\in\mathbb{C}$, set $(x; q, p)_\infty:=\prod_{m, n\geq 0}(1-xq^m p^n)$. For $x\in\mathbb{C}\setminus\{0\}$, set $\Gamma_{q, p}(x):=\frac{(qp x^{-1}; q, p)_\infty}{(x; q, p)_\infty}$ (elliptic gamma function).

Let N be a positive integer, β be a complex number, and p be a complex number satisfying $|p|<1$. The Hamiltonian of the elliptic Calogero-Moser system $H_N^{\text{CM}}(\beta, p)$ is defined by

$$H_N^{\text{CM}}(\beta, p):=\sum_{i=1}^N D_{x_i}^2 - \beta(\beta-1) \sum_{1\leq i\neq j\leq N} E_2(x_i/x_j; p).$$

Then the following fact is known: the function $\Psi_N(x; \beta, p):=\prod_{1\leq i\neq j\leq N} \Theta_p(x_i/x_j)^{\beta/2}$ satisfies

$$H_N^{\text{CM}}(\beta, p)\Psi_N(x; \beta, p)=\{2N\beta D_p + C_N(\beta, p)\}\Psi_N(x; \beta, p), \dots (*)$$

where $C_N(\beta, p)$ is a complex number determined by N, β, p . It is remarkable that the derivative $D_p=p\frac{\partial}{\partial p}$ is in the right hand side of (*). This means that the elliptic Calogero-Moser system has a solution which involves the infinitesimal deformation of the elliptic modulus p .

Let N be a positive integer, q, p be complex numbers satisfying $|q|<1, |p|<1$, and t be a complex number satisfying $t\in\mathbb{C}\setminus\{0\}$. The Hamiltonian of the elliptic Ruijsenaars system $H_N^{\text{R}}(q, t, p)$ is defined by

$$H_N^{\text{R}}(q, t, p):=\sum_{i=1}^N \prod_{j\neq i} \left(\frac{\Theta_p(tx_i/x_j)\Theta_p(qt^{-1}x_i/x_j)}{\Theta_p(x_i/x_j)\Theta_p(qx_i/x_j)} \right)^{\frac{1}{2}} T_{q, x_i},$$

where $T_{q, x}$ is the q -shift operator which is defined by $T_{q, x}f(x)=f(qx)$. Then the function $\Psi_N(x; q, t, p):=\prod_{1\leq i\neq j\leq N} \left(\frac{\Gamma_{q, p}(tx_i/x_j)}{\Gamma_{q, p}(x_i/x_j)} \right)^{1/2}$ satisfies

$$H_N^{\text{R}}(q, t, p)\Psi_N(x; q, t, p)=t^{\frac{-N+1}{2}} \sum_{i=1}^N \prod_{j\neq i} \frac{\Theta_p(tx_i/x_j)}{\Theta_p(x_i/x_j)} \Psi_N(x; q, t, p). \dots (**)$$

It is known that by setting $t=q^\beta$ and by taking the limit $q\rightarrow 1$ appropriately, the equation (**) degenerates to the equation (*). Thus it is probable that the equation (**) contains a certain difference deformation of the elliptic modulus p . By standing the point of view, the author will study the elliptic Ruijsenaars system.