

COHOMOLOGICAL RIGIDITY FOR TORIC FANO MANIFOLDS OF SMALL DIMENSIONS OR LARGE PICARD NUMBERS

AKIHIRO HIGASHITANI, KAZUKI KURIMOTO, AND MIKIYA MASUDA

ABSTRACT. The cohomological rigidity problem for toric manifolds asks whether toric manifolds are diffeomorphic (or homeomorphic) if their integral cohomology rings are isomorphic. Many affirmative partial solutions to the problem have been obtained and no counterexample is known. In this paper, we study the diffeomorphism classification of toric Fano d -folds with $d = 3, 4$ or with Picard number $\geq 2d - 2$. In particular, we show that those manifolds except for two toric Fano 4-folds are diffeomorphic if their integral cohomology rings are isomorphic. The exceptional two toric Fano 4-folds (their ID numbers are 50 and 57 on a list of Øbro) have isomorphic cohomology rings and their total Pontryagin classes are preserved under an isomorphism between their cohomology rings, but we do not know whether they are diffeomorphic or homeomorphic.

1. INTRODUCTION

1.1. Cohomological rigidity problem. As is well-known, integral cohomology ring (as a graded ring) is not a complete invariant to distinguish closed smooth manifolds. However, it becomes a complete invariant if we restrict our concern to a small family \mathcal{F} of closed smooth manifolds. For instance, this is the case if \mathcal{F} is the family of closed surfaces. We say that a family \mathcal{F} of manifolds is *cohomologically rigid* if the integral cohomology rings distinguish the manifolds in \mathcal{F} up to diffeomorphism (or homeomorphism).

A toric variety is a normal complex algebraic variety with an algebraic action of a \mathbb{C}^* -torus having an open dense orbit. It is well known that there is a one-to-one correspondence between toric varieties and a combinatorial object called fans. Therefore, the classification of toric varieties reduces to the classification of fans. A toric variety is not necessarily compact or smooth. A compact smooth toric variety, which we call a *toric manifold*, is well studied. For instance, its cohomology ring and Chern classes are explicitly described in terms of the associated fan. As mentioned above, the classification of toric manifolds *as varieties* reduces to the classification of the associated fans. However, the classification of toric manifolds *as smooth manifolds* is unknown. Motivated by the diffeomorphism classification of a certain family of toric manifolds ([18]), the third author and Dong Youp Suh posed the following naive problem in [19].

Cohomological rigidity problem for toric manifolds. *Are toric manifolds diffeomorphic (or homeomorphic) if their integral cohomology rings are isomorphic as graded rings? Namely, is the family of toric manifolds cohomologically rigid?*

No counterexample to the problem is known and many affirmative partial solutions have been obtained (see [12, 5, 16, 8] and the reference therein for recent accounts of the problem). Among those affirmative solutions, Bott manifolds are well studied. A Bott manifold is a toric manifold associated with a spanning fan of a cross-polytope. It can

2010 *Mathematics Subject Classification.* Primary 14M25; Secondary 57R19, 14J45, 57S15.

Key words and phrases. Cohomological rigidity, toric Fano manifold.

be obtained as the total space of an iterated $\mathbb{C}P^1$ -bundle starting with a point. It is not completely solved that the family of Bott manifolds is cohomologically rigid but the known results are close to the complete solution ([9]). Some partial affirmative solutions are also known for generalized Bott manifolds ([10, 11]) although the known results are far from the complete solution. Here, a generalized Bott manifold is a toric manifold associated with a spanning fan of the direct sum of simplices (see Definition 3.1 for direct sum of polytopes). Similarly to a Bott manifold, a generalized Bott manifold can be obtained as the total space of an iterated $\mathbb{C}P^{n_i}$ -bundle starting with a point, where each n_i can take any positive integer.

In this paper, we study the diffeomorphism classification of toric *Fano* d -folds with $d = 3, 4$ or with Picard numbers $\geq 2d - 2$. Our main result (Theorem 1.1) below provides another affirmative partial solution to the cohomological rigidity problem.

1.2. Smooth Fano d -polytopes and toric Fano d -folds. A *lattice* polytope is a convex polytope $P \subset \mathbb{R}^d$ all of whose vertices lie in the integer lattice \mathbb{Z}^d . We say that $P \subset \mathbb{R}^d$ is a *smooth Fano d -polytope* if it is a full-dimensional lattice polytope containing the origin in its interior such that the set of vertices of every facet forms a \mathbb{Z} -basis of \mathbb{Z}^d . In particular, smooth Fano d -polytopes are simplicial. To a smooth Fano d -polytope $P \subset \mathbb{R}^d$, we can associate a complete nonsingular fan as the *spanning fan* of P , where each i -dimensional cone in the fan is spanned by the vertices of an $(i - 1)$ -dimensional face of P .

It is known that the set of smooth Fano d -polytopes up to unimodular equivalence one-to-one corresponds to the set of toric Fano d -folds up to isomorphism as varieties ([3]). Moreover, it is known that for a fixed d , there are only finitely many smooth Fano d -polytopes up to unimodular equivalence (see [3]). The number of smooth Fano d -polytopes (up to unimodular equivalence) for small values of d is given as follows:

dimension	the number of smooth Fano d -polytopes	proved in
2	5	
3	18	[2, 25]
4	124	[3, 23]
5	866	[20]
6	7622	[20]
7	72256	[20]
8	749892	[20]

Indeed, Øbro ([20]) provides the algorithm (called *SFP algorithm*), which produces a complete list of smooth Fano d -polytopes (up to unimodular equivalence) for a given positive integer d . For $d = 2, 3, 4, 5, 6$, the database of all smooth Fano d -polytopes is open in the following URL:

<http://www.grdb.co.uk/forms/toricsmooth>

Each toric Fano d -fold (or smooth Fano d -polytope) has its ID. For example, the ID number of Hirzebruch surface of degree 0 (resp. 1) is 4 (resp. 3). Toric Fano 3-folds have ID 6–23 and toric Fano 4-folds have ID 24–147, and so on.

One can see that among eighteen toric Fano 3-folds, there are five Bott manifolds (nine generalized Bott manifolds including Bott manifolds), and among one hundred twenty-four toric Fano 4-folds, there are thirteen Bott manifolds (forty-one generalized Bott manifolds including Bott manifolds).

1.3. Results. As explained above, there are only finitely many smooth Fano d -polytopes for a given d and we know their explicit description. This finiteness and explicitness are a great advantage to investigate the cohomological rigidity problem for toric Fano d -folds. In this paper we prove the following.

Theorem 1.1. *The family \mathcal{F} of toric Fano d -folds satisfying one of the following condition:*

- (1) *Picard number $\geq 2d - 2$,*
- (2) *$d = 3$,*
- (3) *$d = 4$ except for ID numbers 50 and 57,*

is cohomologically rigid. Namely, two toric Fano d -folds in the family \mathcal{F} are diffeomorphic if and only if their integral cohomology rings are isomorphic as graded rings.

We actually identify which toric Fano d -folds in the family \mathcal{F} are diffeomorphic (see Tables 1 and 6). Those diffeomorphic toric Fano d -folds are in fact weakly equivariantly diffeomorphic with respect to the restricted actions of the compact subtorus of the \mathbb{C}^* -torus. They also show that the main theorem in [17] is incorrect, see Remark 2.4 (3) for details. Toric Fano 4-folds with ID numbers 50 and 57 have isomorphic integral cohomology rings and their Pontryagin classes are preserved under an isomorphism between their cohomology rings. We do not know whether they are diffeomorphic or homeomorphic, but they are not weakly equivariantly homeomorphic with respect to the restricted actions of the compact subtorus.

The Picard number of a toric Fano d -fold is the number of the vertices of the associated smooth Fano d -polytope minus d . It is known that smooth Fano d -polytopes have at most $3d$ vertices and those with at least $3d - 2$ vertices are classified ([1, 6, 21]). We use this classification result to verify case (1) in Theorem 1.1. As for cases (2) and (3), we use the database of Øbro.

Our approach to the diffeomorphism classification above consists of two directions: one direction is to check an algebraic condition described in terms of fans for two toric manifolds to be homeomorphic or diffeomorphic (Lemma 2.3). The other direction is to use the cohomology rings to distinguish the diffeomorphism classes. It is not difficult to carry out the former direction, but it is quite a task to carry out the latter direction in general.

The cohomology ring $H^*(X)$ of a toric manifold X is the quotient of a polynomial ring in $b_2(X)$ variables by an ideal, where $b_2(X)$ is the rank of $H^2(X)$, i.e. the Picard number of X . In some cases, one can check by an elementary method whether two cohomology rings are isomorphic or not. However, in general, the elementary method requires a formidable computation and does not work well. In order to distinguish our cohomology rings, we pay attention to elements of $H^2(X; R)$ whose k -th power vanish, where we take $R = \mathbb{Z}$, $\mathbb{Z}/2$ or $\mathbb{Z}/3$ and $k = 2, 3$ or 4 . It turns out that they are useful invariants to distinguish our cohomology rings. We often use Gröbner basis to compute those invariants.

Very recently, motivated by McDuff's question on the uniqueness of toric actions on a monotone symplectic manifold, Y. Cho, E. Lee, S. Park and the third author made the following conjecture and verified it for Fano Bott manifolds ([7]).

Conjecture ([7]). *If there is a cohomology ring isomorphism between toric Fano manifolds which preserves their first Chern classes, then they are isomorphic as varieties.*

Based on the classification of cohomology rings of our toric Fano manifolds, we prove

Theorem 1.2. *The conjecture above is true for toric Fano d -folds with $d = 3, 4$ or with Picard number $\geq 2d - 2$.*

1.4. Structure of the paper. In Section 2, we briefly recall the theory of toric varieties and the well-known presentation of the cohomology ring of a toric manifold (Proposition 2.1). We introduce the invariants of cohomology rings used to distinguish our cohomology rings. We also give a lemma (Lemma 2.3) mentioned above to find diffeomorphic or homeomorphic toric manifolds. We recall the notion of Gröbner basis and normal forms. After those preparations, we prove Theorem 1.1. We will verify cases (1), (2), (3) of Theorem 1.1 in Sections 3, 4, 5 respectively. Theorem 1.2 will be proved in Section 6.

Acknowledgements. A. Higashitani was supported in part by JSPS Grant-in-Aid for Scientific Research 18H01134 and M. Masuda was supported in part by JSPS Grant-in-Aid for Scientific Research 16K05152. This work was partly supported by Osaka City University Advanced Mathematical Institute (MEXT Joint Usage/Research Center on Mathematics and Theoretical Physics).

2. PRELIMINARIES

In this section, we recall the well-known presentation of the cohomology ring of a toric manifold (i.e. a compact smooth toric variety) and give a sufficient condition (Lemma 2.3) for two compact smooth toric varieties to be homeomorphic or diffeomorphic. We also introduce naive invariants of cohomology rings used in this paper. We show by an example how to compute those invariants using Gröbner basis.

2.1. Toric manifolds and their cohomology rings. We briefly review the theory of toric varieties and refer the reader to [15] or [22] for details.

A toric variety of complex dimension d is a normal algebraic variety X over the complex numbers \mathbb{C} with an algebraic action of $(\mathbb{C}^*)^d$ having an open dense orbit. A fan in a lattice $N := \text{Hom}(\mathbb{C}^*, (\mathbb{C}^*)^d) (\cong \mathbb{Z}^d)$ is a set of rational strongly convex polyhedral cones in $N \otimes \mathbb{R}$ such that

- (1) each face of a cone in Δ is also a cone in Δ ;
- (2) the intersection of two cones in Δ is a face of each.

The fundamental theorem in the theory of toric varieties says that there is a one-to-one correspondence between toric varieties of complex dimension d and fans in N , and that two toric varieties are isomorphic if and only if the corresponding fans are unimodularly equivalent.

For a fan Δ , we denote the corresponding toric variety by $X(\Delta)$. The toric variety $X(\Delta)$ is compact (or complete) if and only if Δ is complete, i.e. the union of cones in Δ covers the entire space $N \otimes \mathbb{R}$. Moreover, $X(\Delta)$ is smooth (or nonsingular) if and only if Δ is nonsingular, i.e. for any cone σ in Δ , the primitive vectors lying on the 1-dimensional faces of σ form a part of a basis of N .

In the following, we assume that our fan Δ is complete and nonsingular. Let ρ_1, \dots, ρ_m be 1-dimensional cones in Δ . We denote the primitive vector lying on ρ_i by v_i and call it a *primitive ray vector*. We denote the set of all primitive ray vectors by $\mathcal{V}(\Delta)$. The set of all cones in Δ defines an abstract simplicial complex on $[m] = \{1, \dots, m\}$. It is called the *underlying simplicial complex* of Δ and denoted by $\mathcal{K}(\Delta)$. Indeed, a subset I of $[m]$ is a member of $\mathcal{K}(\Delta)$ if and only if v_i 's for $i \in I$ span a cone in Δ . Since Δ can be recovered from two data $\mathcal{K}(\Delta)$ and $\mathcal{V}(\Delta)$, we may think of Δ as the pair $(\mathcal{K}(\Delta), \mathcal{V}(\Delta))$.

Each 1-dimensional cone ρ_i in the fan Δ corresponds to an invariant divisor X_i of $X(\Delta)$ and we denote the Poincaré dual to X_i by x_i . Since X_i is of real codimension two, x_i lies in $H^2(X(\Delta))$. With this understanding, we have the following well-known presentation of the cohomology ring of a toric manifold. Throughout this paper, we will use it without mentioning it.

Proposition 2.1 ([4, Theorem 5.3.1]). *The cohomology ring of a toric manifold $X(\Delta)$ can be described as follows:*

$$H^*(X(\Delta)) = \mathbb{Z}[x_1, \dots, x_m] / \mathcal{J}$$

where $\deg x_i = 2$ for $i = 1, \dots, m$ and \mathcal{J} is the ideal generated by all

- (i) $x_{i_1} \cdots x_{i_k}$ for $\{i_1, \dots, i_k\} \notin \mathcal{K}(\Delta)$;
- (ii) $\sum_{j=1}^m u(v_j)x_j$ for $u \in \text{Hom}(N, \mathbb{Z})$.

Remark 2.2. (1) The total Chern class and the total Pontryagin class of $X(\Delta)$ are respectively given by

$$c(X(\Delta)) = \prod_{i=1}^m (1 + x_i), \quad p(X(\Delta)) = \prod_{i=1}^m (1 + x_i^2).$$

(2) Since the fan Δ is complete and nonsingular, one can eliminate d variables among x_1, \dots, x_m using the linear relations (ii), so $H^*(X(\Delta))$ is actually the quotient of a polynomial ring in $m - d$ variables by an ideal, where $m - d$ agrees with the rank of $H^2(X(\Delta))$ because $k \geq 2$ in (i).

2.2. Diffeomorphism lemma. In this subsection, we give a sufficient condition for toric manifolds to be homeomorphic or diffeomorphic. By definition, a toric manifold $X(\Delta)$ of complex dimension d has an algebraic action of $(\mathbb{C}^*)^d$. Let S^1 be the unit circle group of \mathbb{C} . It is known that the orbit space Q of $X(\Delta)$ by the restricted action of $(S^1)^d$ is a d -dimensional manifold with corners such that all faces (even Q itself) are contractible. The dual face poset of Q defines a simplicial complex which agrees with the simplicial complex $\mathcal{K}(\Delta)$, to be more precise, if Q_1, \dots, Q_m are the facets of Q , then $Q_I := \bigcap_{i \in I} Q_i \neq \emptyset$ for $I \subset [m]$ if and only if $I \in \mathcal{K}(\Delta)$. The orbit space Q is often a simple polytope. In fact, this is the case when $X(\Delta)$ is projective, more generally when $\mathcal{K}(\Delta)$ is the boundary complex of a simplicial polytope.

Remember that $\mathcal{V}(\Delta) = \{v_1, \dots, v_m\}$ is the set of primitive ray vectors in Δ . Since $N = \text{Hom}(\mathbb{C}^*, (\mathbb{C}^*)^d) = \text{Hom}(S^1, (S^1)^d)$, we may regard $v_i \in N$ as a homomorphism from S^1 to $(S^1)^d$. We note that Q is the disjoint union of the interior part $\text{Int } Q_I$ of Q_I over $I \in \mathcal{K}(\Delta)$, where we allow $I = \emptyset$ and $Q_\emptyset = Q$. We consider the quotient space

$$X(Q, \mathcal{V}(\Delta)) := Q \times (S^1)^d / \sim \tag{2.1}$$

where $(x, g) \sim (y, h)$ if and only if $x = y \in \text{Int } Q_I$ and gh^{-1} belongs to the subtorus generated by circle subgroups $v_i(S^1)$ of $(S^1)^d$ for $i \in I$. The space $X(Q, \mathcal{V}(\Delta))$ is called the *canonical model* of $X(\Delta)$ because $X(Q, \mathcal{V}(\Delta))$ is homeomorphic to $X(\Delta)$ ([14], [4, Chapter 7]).

Lemma 2.3. *Let Δ and Δ' be complete nonsingular fans with the same underlying simplicial complex, i.e. $\mathcal{K}(\Delta) = \mathcal{K}(\Delta')$. If the sets of primitive ray vectors $\mathcal{V}(\Delta) = \{v_1, \dots, v_m\}$ and $\mathcal{V}(\Delta') = \{v'_1, \dots, v'_m\}$ agree up to sign (which means $v_i = \pm v'_i$ for each $i = 1, \dots, m$), then $X(\Delta)$ and $X(\Delta')$ are homeomorphic. Moreover, if $\mathcal{K}(\Delta) = \mathcal{K}(\Delta')$ is the boundary*

complex of a simplicial polytope, then “homeomorphic” above can be improved to “diffeomorphic”.

Proof. As remarked above, $X(\Delta)$ is homeomorphic to the canonical model $X(Q, \mathcal{V}(\Delta))$. We see from the construction that the canonical model does not depend on the signs of v_i 's, so the former statement follows. If $\mathcal{K}(\Delta) = \mathcal{K}(\Delta')$ is the boundary complex of a simplicial polytope, then the orbit space Q is a simple polytope; so the latter statement follows from [13, Proposition 6.4 (iii)]. \square

Remark 2.4. (1) The multiplication by $(S^1)^d$ on the second factor of $X(Q, \mathcal{V}(\Delta))$ descends to an action of $(S^1)^d$ on $X(Q, \mathcal{V}(\Delta))$ and Lemma 2.3 holds in this equivariant setting, see [4, Chapter 7].

(2) If two complete nonsingular fans are unimodularly equivalent, then the associated two toric manifolds are not only isomorphic but also weakly equivariantly isomorphic with respect to the actions of $(\mathbb{C}^*)^d$ as is known.

(3) It is proved in the paper [17] by the third author that if the equivariant cohomology rings of two toric manifolds X and X' are isomorphic as algebras over $H^*(BT)$, where T is the \mathbb{C}^* -torus acting on X and X' , then $v_i = \pm v'_i$ (i.e. the condition in Lemma 2.3 is satisfied). After that, the author claimed that it implies that X and X' are isomorphic as varieties but this is incorrect. The mistake occurs at the very end of the proof of Theorem 1 in [17] and the author thanks Hiraku Abe for pointing out the mistake. Indeed, we will see in this paper that there are toric Fano manifolds which satisfy the condition in Lemma 2.3 but they are not isomorphic as varieties. The author also thanks Hiroshi Sato for providing such an example. In order to correct the theorem, we need to take equivariant first Chern class into account. Namely, if the equivariant cohomology algebra isomorphism between X and X' preserves their equivariant first Chern classes (those are $\sum_{i=1}^m \tau_i$ and $\sum_{i=1}^m \tau'_i$ in [17]), then we can conclude $v_i = v'_i$ for every i so that X and X' are isomorphic as varieties.

2.3. Invariants of a cohomology ring. Let Δ be a complete nonsingular fan of dimension d . It is well-known that the number of i -dimensional cones in Δ coincides with the $2i$ -th Betti number of the toric manifold $X(\Delta)$ for $0 \leq i \leq d-1$. Therefore, when Δ is the spanning fan of a simplicial d -polytope P , the face numbers of P are cohomology invariants. Namely, if toric manifolds associated with simplicial polytopes P and P' have isomorphic cohomology rings, then the face numbers of P and P' coincide. Especially, the number of vertices of P and the number of facets of P are cohomology invariants. We will use this fact without mentioning it throughout this paper.

Betti numbers are invariants of a cohomology ring but they depend only on its additive structure. We now introduce invariants of a cohomology ring, which depend on its ring structure.

Definition 2.5 (*s.v.e., c.v.e., k-v.e.*). Let $k \geq 2$ and $R = \mathbb{Z}$ or \mathbb{Z}/p where p is a prime number. We say that a nonzero element of $H^2(X) \otimes R$ is *k-v.e.* over R if it is primitive and its k -th power vanishes in $H^*(X) \otimes R$. When $R = \mathbb{Z}$, the word “over \mathbb{Z} ” will be omitted. Also, 2-v.e. will be called *s.v.e.* (square vanishing element) and 3-v.e. will be called *c.v.e.* (cube vanishing element).

Definition 2.6 (*maximal basis number*). Let V be the set of all s.v.e. of $H^*(X)$ and we consider

$$\mathcal{B} := \{S \subset V \mid S \text{ is a part of a } \mathbb{Z}\text{-basis of } H^2(X)\}.$$

Clearly there exists an $S_{\max} \in \mathcal{B}$ such that

$$|S| \leq |S_{\max}| \quad \text{for } \forall S \in \mathcal{B}.$$

We call $|S_{\max}|$ the *maximal basis number* of $H^*(X)$.

The number of k -v.e. over R and the maximal basis number are invariants of $H^*(X)$. Note that there may exist infinitely many s.v.e. although the maximal basis number is finite.

Example 2.7. We compute s.v.e. and the maximal basis numbers of $H^*(F_0)$ and $H^*(F_1)$, where F_a is the Hirzebruch surface of degree a . As is well-known, we have

$$H^*(F_0) \cong \mathbb{Z}[x, y]/(x^2, y^2) \quad \text{and} \quad H^*(F_1) \cong \mathbb{Z}[x, y]/(x^2, y(y-x)).$$

Then we easily obtain the following table.

	s.v.e. (up to sign)	maximal basis number
F_0	x, y	2
F_1	$x, x - 2y$	1

The number of s.v.e. are the same but the maximal basis numbers are different. Thus, $H^*(F_0)$ and $H^*(F_1)$ are not isomorphic to each other.

2.4. Gröbner basis and normal forms. For the computation of k -v.e. of $H^*(M)$, we recall what Gröbner basis is. We refer the reader to [24] for the introduction to Gröbner basis. The terminologies not defined in this section for Gröbner basis can be found there.

Let $S = \mathbb{k}[x_1, \dots, x_m]$ be a polynomial ring with m variables over a field \mathbb{k} and let \mathcal{M} be the set of all monomials in S . We say that a total order $<$ in \mathcal{M} a *monomial order* on S if it satisfies that

- $1 < u$ for any $u \in \mathcal{M}$ with $u \neq 1$, and
- $uw < vw$ holds for any $u, v, w \in \mathcal{M}$ with $u < v$.

Fix a monomial order $<$ on S . Given a polynomial $f \in S$, we call the leading monomial with respect to $<$ appearing in f the *initial monomial*, denoted by $\text{in}_{<}(f)$. Given an ideal $I \subset S$, the ideal generated by the initial monomials f in I is called the *initial ideal* of I , denoted by $\text{in}_{<}(I)$. Namely,

$$\text{in}_{<}(I) = (\text{in}_{<}(f) \mid f \in I).$$

For a system of generator $\{g_1, \dots, g_s\}$ of an ideal I , it is not necessarily the case that $\text{in}_{<}(I) = (\text{in}_{<}(g_1), \dots, \text{in}_{<}(g_s))$ holds. We say that $\{g_1, \dots, g_s\}$ is a *Gröbner basis for I with respect to a monomial order $<$* if this equality holds. Even if a system of generator \mathcal{G} of an ideal is not a Gröbner basis for an ideal, there is an algorithm, so called *Buchberger algorithm*, to compute its Gröbner basis from \mathcal{G} by appending some additional generators to \mathcal{G} .

Let $I \subset S$ be an ideal and let $\{g_1, \dots, g_s\} \subset I$ be a system of generator of I . For $f \in S$ which is not equal to 0, we can get an equation

$$f = f_1g_1 + \dots + f_sg_s + f'$$

satisfying the following:

- $u \notin (\text{in}_{<}(g_1), \dots, \text{in}_{<}(g_s))$ for all monomials u appearing in f' if $f' \neq 0$, and
- $\text{in}_{<}(f_i g_i) \leq \text{in}_{<}(f)$ if $f_i \neq 0$.

We call f' a *normal form* of f and write $\text{NF}(f)$. It is known that $\text{NF}(f)$ is well-defined if $\{g_1, \dots, g_s\}$ is a Gröbner basis for I .

Once we get a Gröbner basis of an ideal I , we have many advantages. One of such advantages is the following proposition:

Proposition 2.8. *Let $I \subset S$ be an ideal and let $\{g_1, \dots, g_s\}$ be a Gröbner basis for I with respect to a monomial order $<$. Given a polynomial $f \in S$, we have the following:*

$$f \in I \iff \text{NF}(f) = 0.$$

Example 2.9. Let $X = X_{12}$ be the toric Fano manifold corresponding to ID number 12, which will appear in Section 4, and let P be the associated smooth Fano 3-polytope. Then P has six vertices.

Let us consider the cohomology ring $H^*(X)$, and demonstrate how to compute s.v.e. of this ring. According to the database, the vertices of P are as follows:

$$v_1 = (1, 0, 0), v_2 = (0, 1, 0), v_3 = (0, 0, 1), v_4 = (-1, 0, 1), v_5 = (0, 1, -1), v_6 = (0, -1, 0).$$

1) First, we compute the defining ideal \mathcal{I} of the cohomology ring. By Proposition 2.1, we can compute \mathcal{I} as follows:

$$H^*(X) \cong \mathbb{Z}[x_1, \dots, x_6]/(I_X + J_X), \text{ where}$$

$$I_X = (x_1x_4, x_2x_6, x_3x_5) \text{ and } J_X = (x_1 - x_4, x_2 + x_5 - x_6, x_3 + x_4 - x_5).$$

Note that I_X (resp. J_X) corresponds to (i) (resp. (ii)). So, we obtain that

$$\begin{aligned} H^*(X) &\cong \mathbb{Z}[x_1, \dots, x_6]/((x_1x_4, x_2x_6, x_3x_5) + (x_1 - x_4, x_2 + x_5 - x_6, x_3 + x_4 - x_5)) \\ &\cong \mathbb{Z}[x, y, z]/(x^2, (z - y)z, (y - x)y). \end{aligned}$$

Note that we apply the change of variables $x_4 = x$, $x_5 = y$, $x_6 = z$, and $x_1 = x$, $x_2 = z - y$ and $x_3 = y - x$.

2) Next, we compute a Gröbner basis for the ideal $\mathcal{I} = (x^2, z(z - y), y(y - x))$. As a monomial order, let $<$ be the graded lexicographic order induced by $x < y < z$. Then we can see that $\{x^2, z(z - y), y(y - x)\}$ is a Gröbner basis for \mathcal{I} with respect to $<$.

3) Finally, we compute s.v.e. of $H^*(X)$. Let $f_1 = x^2$, $f_2 = z^2 - yz$ and $f_3 = y^2 - xy$.

Let $f = ax + by + cz$. Then the normal form of f^2 can be computed as follows:

$$\begin{aligned} f^2 &= a^2x^2 + b^2y^2 + c^2z^2 + 2abxy + 2acxz + 2bcyz \\ &= a^2f_1 + c^2f_2 + b^2f_3 + (2a + b)bx y + 2acxz + (2b + c)cyz. \end{aligned}$$

Hence, $\text{NF}(f^2) = (2a + b)bx y + 2acxz + (2b + c)cyz$. Thus, we can see that $\text{NF}(f^2) = 0$ if and only if

$$\begin{cases} (2a + b)b = 0, \\ 2ac = 0, \\ (2b + c)c = 0. \end{cases}$$

When $a = 0$, we have $b = c = 0$. When $a \neq 0$, we have $c = 0$ and $(2a + b)b = 0$, i.e., $b = 0$ or $b = -2a$. Hence, we conclude that $\text{NF}(f^2) = 0$ if and only if $f = ax$ or $f = ax - 2ay$. Therefore, s.v.e. of $H^*(X)$ are x and $x - 2y$, and its maximal basis number is 1 since x and $x - 2y$ cannot form a \mathbb{Z} -basis.

Example 2.10. Let us consider the cohomology ring $H^*(X_{24})$, which will appear in Section 5, and demonstrate how to compute s.v.e. of this ring. Let $X = X_{24}$ and let P

be the associated smooth Fano 4-polytope. Then P has seven vertices. According to the database, the vertices of P are as follows:

$$\begin{aligned} v_1 &= (1, 0, 0, 0), & v_2 &= (0, 1, 0, 0), & v_3 &= (0, 0, 1, 0), & v_4 &= (0, 0, 0, 1), \\ v_5 &= (-1, -1, -1, 3), & v_6 &= (0, 0, 1, -1), & v_7 &= (0, 0, 0, -1). \end{aligned}$$

1) First, we compute the defining ideal \mathcal{I} of the cohomology ring. By Proposition 2.1, we can compute \mathcal{I} as follows:

$$\begin{aligned} H^*(X) &\cong \mathbb{Z}[x_1, \dots, x_8]/(I_X + J_X), \text{ where} \\ I_X &= (x_1x_2x_3x_5, x_3x_7, x_4x_7, x_4x_6, x_1x_2x_5x_6) \text{ and} \\ J_X &= (x_1 - x_5, x_2 - x_5, x_3 - x_5 + x_6, x_4 + 3x_5 - x_6 - x_7), \\ &\cong \mathbb{Z}[x, y, z]/(x^3(x-y), (x-y)z, (-3x+y+z)z, (-3x+y+z)y, x^3y) \\ &= \mathbb{Z}[x, y, z]/(x^4, (x-y)z, (-2y+z)z, (-2x+y)y, x^3y). \end{aligned}$$

Note that we apply the change of variables $x_5 = x$, $x_6 = y$, $x_7 = z$, and $x_1 = x_2 = x$, $x_3 = x - y$ and $x_4 = -3x + y + z$.

2) Next, we compute a Gröbner basis for the ideal

$$\mathcal{I} = (x^4, (x-y)z, (-2y+z)z, (-3x+y+z)y, x^3y).$$

As a monomial order, let $<$ be the graded lexicographic order induced by $x < y < z$. A Gröbner basis for I with respect to a monomial order $<$ is

$$\{x^4, (x-y)z, (-2y+z)z, (-2x+y)y, x^3y, x^2z\}. \quad (2.2)$$

Note that the generator itself does not become a Gröbner basis since

$$\mathcal{I} \ni (x-y) \cdot (x-y)z - z \cdot (-2x+y)y = x^2z \notin \text{in}_{<}(\mathcal{I}),$$

but it follows from Buchberger algorithm that (2.2) is a Gröbner basis for \mathcal{I} with respect to $<$.

3) Finally, we compute s.v.e. of $H^*(X)$. Let $f = ax + by + cz$. Then $\text{NF}(f^2) = a^2x^2 + 2b(a+b)xy + 2c(a+b+c)xz$. Therefore, we conclude that $\text{NF}(f^2) = 0$ if and only if $a = b = c = 0$. This means that $H^*(X)$ has no s.v.e.

3. THE CASE WITH LARGE PICARD NUMBER

This section is devoted to verifying the case (1) in Theorem 1.1. The Picard number of a smooth Fano d -fold associated with a smooth Fano d -polytope P is the number of vertices of P minus d . On the other hand, smooth Fano d -polytopes are known to have at most $3d$ vertices and those with $3d$, $3d - 1$ or $3d - 2$ vertices are classified. We prepare some terminology to state those results.

Definition 3.1 (*direct sum*). Let $P \subset \mathbb{R}^d$ and $Q \subset \mathbb{R}^e$ be polytopes. Then

$$P \oplus Q = \text{conv}(P \times \{\mathbf{0}_e\} \cup \{\mathbf{0}_d\} \times Q) \subset \mathbb{R}^{d+e},$$

where $\mathbf{0}_d$ (resp. $\mathbf{0}_e$) denotes the origin of \mathbb{R}^d (resp. \mathbb{R}^e), is called the *direct sum* of P and Q . The direct sum is also called the *free sum* of P and Q .

Definition 3.2 (*skew bipyramid*). Let $P \subset \mathbb{R}^d$ be a polytope. Then we call a polytope $B \subset \mathbb{R}^{d+1}$ a *skew bipyramid* (or simply *bipyramid*) over P if P is contained in affine hyperplane H such that there are two vertices v and w of B which lie on either side of H such that $B = \text{conv}(\{v, w\} \cup P)$ and the line segment $[v, w]$ meets P in its (relative) interior.

Throughout this section, e_1, \dots, e_d will denote the standard basis of \mathbb{Z}^d and P_6 (resp. P_5) will denote the hexagon (resp. pentagon) with vertices

$$\pm e_1, \pm e_2, \pm(e_1 - e_2) \quad (\text{resp. } e_1, \pm e_2, \pm(e_1 - e_2)).$$

The classification results on smooth Fano d -polytopes with $3d$, $3d - 1$ or $3d - 2$ vertices are as follows.

Theorem 3.3 (Casagrande [6]). *A smooth Fano d -polytope P has at most $3d$ vertices. If it does have exactly $3d$ vertices, then d is even and P is unimodularly equivalent to $P_6^{\oplus \frac{d}{2}}$.*

Theorem 3.4 (Øbro [21]). *Let P be a smooth Fano d -polytope with exactly $3d - 1$ vertices. If d is even, then P is unimodularly equivalent to $P_5 \oplus P_6^{\oplus \frac{d-2}{2}}$. If d is odd, then P is unimodularly equivalent to a bipyramid over $P_6^{\oplus \frac{d-1}{2}}$.*

Theorem 3.5 (Assarf-Joswig-Paffenholz [1]). *Let P be a smooth Fano d -polytope with exactly $3d - 2$ vertices. If d is even, then P is unimodularly equivalent to*

$$\begin{cases} (1) \text{ a double bipyramid over } P_6^{\oplus \frac{d-2}{2}} & \text{or} \\ (2) P_5 \oplus P_5 \oplus P_6^{\oplus \frac{d-4}{2}} & \text{or} \\ (3) DP(4) \oplus P_6^{\oplus \frac{d-4}{2}}, \end{cases}$$

where $DP(4)$ is the convex hull of 10 vertices $\pm e_1, \pm e_2, \pm e_3, \pm e_4, \pm(e_1 + e_2 + e_3 + e_4)$ in \mathbb{R}^4 . If d is odd, then P is unimodularly equivalent to a bipyramid over $P_5 \oplus P_6^{\oplus \frac{d-3}{2}}$.

Remark 3.6. The polytopes in (1), (2), (3) in Theorem 3.5 have different face numbers. Indeed, their facet numbers are respectively $24 \cdot 6^{\frac{d-4}{2}}$, $25 \cdot 6^{\frac{d-4}{2}}$, $30 \cdot 6^{\frac{d-4}{2}}$.

3.1. Cohomology of toric Fano 2-folds associated to P_6 and P_5 . Let X_{P_6} be the toric Fano 2-fold associated to P_6 . We number the vertices of P_6

$$e_1, e_2, -e_1 + e_2, -e_1, -e_2, e_1 - e_2$$

from 1 to 6 and denote the corresponding elements in $H^2(X_{P_6})$ by μ_1, \dots, μ_6 . Then we have

$$\begin{aligned} H^*(X_{P_6}) &\cong \mathbb{Z}[\mu_1, \mu_2, \mu_3, \mu_4, \mu_5, \mu_6] / ((\mu_1\mu_3, \mu_1\mu_4, \mu_1\mu_5, \mu_2\mu_4, \mu_2\mu_5, \mu_2\mu_6, \mu_3\mu_5, \mu_3\mu_6, \mu_4\mu_6) \\ &\quad + (\mu_1 - \mu_3 - \mu_4 + \mu_6, \mu_2 + \mu_3 - \mu_5 - \mu_6)) \\ &\cong \mathbb{Z}[x, y, z, w] / (x(x+y), y(x+y), z(y-w), y(x-z), z(z+w), w(z+w), xz, xw, yw), \end{aligned} \tag{3.1}$$

where $x = \mu_3$, $y = \mu_4$, $z = \mu_5$ and $w = \mu_6$.

Lemma 3.7. *The maximal basis number of $H^*(X_{P_6})$ is 3.*

Proof. It follows from (3.1) that any element of $H^2(X_{P_6})$ is of the form $ax + by + cz + dw$ with integers a, b, c, d and an elementary computation shows that

$$(ax + by + cz + dw)^2 = (a^2 - 2ab + b^2 - 2bc + c^2 - 2cd + d^2)w^2.$$

Therefore, the s.v.e of $H^*(X_{P_6})$ are primitive elements in the set

$$\{ax + by + cz + dw \mid (a - b)^2 + (c - d)^2 = 2bc\}. \tag{3.2}$$

In particular, $x + y$, $y + z$ and $z + w$ are s.v.e. and since they form a part of a \mathbb{Z} -basis of $H^2(X_{P_6})$, the maximal basis number of $H^*(X_{P_6})$ is at least 3.

On the other hand, it follows from (3.2) that s.v.e. of $H^*(X_{P_6})$ over $\mathbb{Z}/2$ are given by

$$\{ax + by + cz + dw \mid a + b + c + d = 0\},$$

where a, b, c, d are regarded as elements of $\mathbb{Z}/2$. Therefore, the dimension of $H^2(X_{P_6}) \otimes \mathbb{Z}/2$ is 3, which implies that the maximal basis number of $H^*(X_{P_6})$ is at most 3, proving the lemma. \square

Lemma 3.8. *For any s.v.e. f of $H^*(X_{P_6})$, there exist infinitely many s.v.e. of $H^*(X_{P_6})$ such that f together with the s.v.e. does not form a part of a \mathbb{Z} -basis of $H^2(X_{P_6})$.*

Proof. As observed in (3.2), any s.v.e. $ax + by + cz + dw$ of $H^*(X_{P_6})$ must satisfy the condition

$$(a - b)^2 + (c - d)^2 = 2bc.$$

Therefore, the parity of $a - b$ and $c - d$ must be the same. Moreover, at least one of a, b, c, d must be odd because the s.v.e. $ax + by + cz + dw$ is primitive. In fact, one can see from the identity above that the parity of (a, b, c, d) must be one of the following up to symmetry:

- (i) (even, even, odd, odd); (ii) (even, odd, odd, even).

Since any two elements with the same parity do not form a part of a \mathbb{Z} -basis of $H^2(X_{P_6})$, it suffices to show that there are infinitely many s.v.e. in each of (i) and (ii) above and here are examples of infinitely many s.v.e. in each case:

- (i) $(2k(k + 1), 2k^2, 1, 1)$, (ii) $(2(k + 1)^2, 2k(k + 1) + 1, 1, 0)$,

where k is any integer. \square

Let X_{P_5} be the toric Fano 2-fold associated to P_5 . We number the vertices of P_5

$$e_1, e_2, -e_1 + e_2, -e_2, e_1 - e_2$$

from 1 to 5 and denote the corresponding elements in $H^2(X_{P_5})$ by μ_1, \dots, μ_5 . Then we have

$$\begin{aligned} H^*(X_{P_5}) &\cong \mathbb{Z}[\mu_1, \mu_2, \mu_3, \mu_4, \mu_5] / ((\mu_1\mu_3, \mu_1\mu_4, \mu_2\mu_4, \mu_2\mu_5, \mu_3\mu_5) \\ &\quad + (\mu_1 - \mu_3 + \mu_5, \mu_2 + \mu_3 - \mu_4 - \mu_5)) \\ &\cong \mathbb{Z}[x, y, z] / (x^2, y(x - z), y^2, z(y + z), xz), \end{aligned} \tag{3.3}$$

where $x = \mu_3$, $y = \mu_4$ and $z = \mu_5$.

Lemma 3.9. *The maximal basis number of $H^*(X_{P_5})$ is 2.*

Proof. The proof is essentially the same as in Lemma 3.7. It follows from (3.3) that any element of $H^2(X_{P_5})$ is of the form $ax + by + cz$ with integers a, b, c and an elementary computation shows that

$$(ax + by + cz)^2 = (-2ab - 2bc + c^2)z^2.$$

Therefore, s.v.e of $H^*(X_{P_5})$ are primitive elements in the set

$$\{ax + by + cz \mid c^2 = 2b(a + c)\}. \tag{3.4}$$

In particular, x and y are s.v.e. and since they form a part of \mathbb{Z} -basis, the maximal basis number of s.v.e. of $H^*(X_{P_5})$ is at least 2.

On the other hand, (3.4) shows that c must be even. This implies that the basis maximal number must be at most 2, proving the lemma. \square

Lemma 3.10. *Let $A = \mathbb{Z}[x_1, \dots, x_m]/\mathcal{I}_A$ and $B = \mathbb{Z}[y_1, \dots, y_n]/\mathcal{I}_B$. If f is an s.v.e. of $A \otimes B$, then $f \in A$ or $f \in B$. In particular, the maximal basis number behaves additively with respect to tensor products.*

Proof. Let $f = \sum_{i=1}^m a_i x_i + \sum_{j=1}^n b_j y_j$. Then,

$$\begin{aligned} f^2 &= \left(\sum_{i=1}^m a_i x_i \right)^2 + \left(\sum_{j=1}^n b_j y_j \right)^2 + 2 \left(\sum_{i=1}^m a_i x_i \right) \left(\sum_{j=1}^n b_j y_j \right) \\ &= \left(\sum_{i=1}^m a_i x_i \right)^2 + \left(\sum_{j=1}^n b_j y_j \right)^2 + 2 \left(\sum_{i=1}^m \sum_{j=1}^n a_i b_j x_i y_j \right) \end{aligned}$$

Thus, we have $a_i b_j = 0$ for any i and j , i.e. we have either $a_i = 0$ for any i or $b_j = 0$ for any j . \square

Under these preparations, we start to prove Theorem 1.1 for the case (1). There are following five cases according to the number of vertices $V(P)$ and the parity of the dimension d for smooth Fano d -polytopes P :

- (1) $V(P) = 3d$ (in this case d must be even),
- (2) $V(P) = 3d - 1$ and d is odd,
- (3) $V(P) = 3d - 1$ and d is even,
- (4) $V(P) = 3d - 2$ and d is odd,
- (5) $V(P) = 3d - 2$ and d is even.

In (1) and (3) above, there is only one smooth Fano d -polytope by Theorems 3.3 and 3.4, so it suffices to treat the remaining three cases. In the following, we may assume $d \geq 3$ since it is known (and easy to see) that smooth toric d -folds are distinguished by their cohomology rings when $d \leq 2$.

3.2. The case where $V(P) = 3d - 1$ with d odd ≥ 3 . By Theorem 3.4, there are two P 's up to unimodular equivalence and their vertices are as follows:

$$e_1, \quad -e_1, \quad \pm e_2, \pm e_3, \pm(e_2 - e_3), \dots, \pm e_{d-1}, \pm e_d, \pm(e_{d-1} - e_d), \quad (Y_1^d)$$

$$e_1, -e_1 + e_2, \pm e_2, \pm e_3, \pm(e_2 - e_3), \dots, \pm e_{d-1}, \pm e_d, \pm(e_{d-1} - e_d). \quad (Y_2^d)$$

where the tags denote the corresponding toric Fano d -folds. In each case, the first two vertices correspond to the apices of the bipyramid and $\{\pm e_{2k}, \pm e_{2k+1}, \pm(e_{2k} - e_{2k+1})\}$ ($1 \leq k \leq \frac{d-1}{2}$) forms the hexagon P_6 . One can see from the above that

$$H^*(Y_1^d) = \mathbb{Z}[x]/(x^2) \otimes H^*(X_{P_6})^{\otimes \frac{d-1}{2}}, \quad H^*(Y_2^d) = H^*(Y_2^3) \otimes H^*(X_{P_6})^{\otimes \frac{d-3}{2}}$$

We number the eight vertices

$$e_2, \quad e_3, \quad -e_2 + e_3, \quad -e_2, \quad -e_3, \quad e_2 - e_3, \quad e_1, \quad -e_1 + e_2$$

in (Y_2^3) above from 1 to 8 and denote the corresponding elements in $H^2(Y_2^3)$ by μ_1, \dots, μ_8 . Then we have

$$\begin{aligned} H^*(Y_2^3) &\cong \mathbb{Z}[\mu_1, \mu_2, \mu_3, \mu_4, \mu_5, \mu_6, \mu_7, \mu_8] / ((\mu_1\mu_3, \mu_1\mu_4, \mu_1\mu_5, \mu_2\mu_4, \mu_2\mu_5, \mu_2\mu_6, \mu_3\mu_5, \mu_3\mu_6, \mu_4\mu_6, \mu_7\mu_8) \\ &\quad + (\mu_1 - \mu_3 - \mu_4 + \mu_6 + \mu_8, \mu_2 + \mu_3 - \mu_5 - \mu_6, \mu_7 - \mu_8)) \\ &\cong \mathbb{Z}[x, y, z, w, v] / (x(x + y - v), y(x + y - v), z(y - w - v), \\ &\quad y(x - z), z(z + w), w(z + w), xz, xw, yw, v^2), \end{aligned}$$

where $x = \mu_3$, $y = \mu_4$, $z = \mu_5$, $w = \mu_6$ and $v = \mu_8$. One can check that

$$\text{s.v.e. of } H^*(Y_2^3) = \{2y + 2z - v, 2x + 2y - v, z + w, v\} \text{ (up to sign),} \quad (3.5)$$

and hence the maximal basis number of $H^*(Y_2^3)$ is 2. Therefore, it follows from Lemma 3.7 and Lemma 3.10 that the maximal basis numbers of $H^*(Y_1^d)$ and $H^*(Y_2^d)$ are as in the following table, so $H^*(Y_1^d)$ and $H^*(Y_2^d)$ are not isomorphic to each other.

Ring	Maximal basis number
$H^*(Y_1^d)$	$1 + 3 \cdot \frac{d-1}{2}$
$H^*(Y_2^d)$	$-1 + 3 \cdot \frac{d-1}{2}$

3.3. The case where $V(P) = 3d - 2$ with d odd ≥ 3 . We first treat the case where $d = 3$. By Theorem 3.5, the vertices of P are one of the following:

$$e_1, \quad -e_1, \quad e_2, \pm e_3, \pm(e_2 - e_3), \quad (Z_1)$$

$$e_1, \quad -e_1 + e_2, \quad e_2, \pm e_3, \pm(e_2 - e_3), \quad (Z_2)$$

$$e_1, \quad -e_1 + e_3, \quad e_2, \pm e_3, \pm(e_2 - e_3), \quad (Z_3)$$

$$e_1, \quad -e_1 - e_3, \quad e_2, \pm e_3, \pm(e_2 - e_3). \quad (Z_4)$$

In each case, $\{e_2, \pm e_3, \pm(e_2 - e_3)\}$ forms P_5 and the first two vertices correspond to the apices of the bipyramid over P_5 .

We claim that Z_3 and Z_4 are diffeomorphic. Indeed, the vertices in Z_3 are unimodularly equivalent to

$$-e_1, \quad e_1 + e_3, \quad e_2, \pm e_3, \pm(e_2 - e_3)$$

through an automorphism $(x_1, x_2, x_3) \rightarrow (-x_1, x_2, x_3)$ of \mathbb{Z}^3 , and these vectors agree with the vectors in (Z_4) up to sign. Therefore, Z_3 and Z_4 are diffeomorphic by Lemma 2.3.

We shall observe that the cohomology rings of Z_1, Z_2, Z_3 are not isomorphic to each other. Clearly $Z_1 = \mathbb{C}P^1 \times X_{P_5}$ and hence

$$H^*(Z_1) \cong \mathbb{Z}[x]/(x^2) \otimes H^*(X_{P_5}).$$

To describe the cohomology rings of the remaining ones, we number the seven vertices

$$e_2, \quad e_3, \quad -e_2 + e_3, \quad -e_3, \quad e_2 - e_3, \quad e_1, \quad -e_1 + *$$

from 1 to 7, where $*$ = e_2 or $\pm e_3$. We denote the corresponding elements in $H^2(Z_i)$ for $i = 2, 3$ by μ_1, \dots, μ_7 and set

$$x = \mu_3, \quad y = \mu_4, \quad z = \mu_5, \quad w = \mu_7.$$

Then we have

$$\begin{aligned}
H^*(Z_2) &\cong \mathbb{Z}[\mu_1, \mu_2, \mu_3, \mu_4, \mu_5, \mu_6, \mu_7] / ((\mu_1\mu_3, \mu_1\mu_4, \mu_2\mu_4, \mu_2\mu_5, \mu_3\mu_5, \mu_6\mu_7) \\
&\quad + (\mu_1 - \mu_3 + \mu_5 + \mu_7, \mu_2 + \mu_3 - \mu_4 - \mu_5, \mu_6 - \mu_7)) \\
&\cong \mathbb{Z}[x, y, z, w] / (x(x-w), y(x-z-w), y(y-w), z(y+z), xz, w^2); \\
H^*(Z_3) &\cong \mathbb{Z}[\mu_1, \mu_2, \mu_3, \mu_4, \mu_5, \mu_6, \mu_7] / ((\mu_1\mu_3, \mu_1\mu_4, \mu_2\mu_4, \mu_2\mu_5, \mu_3\mu_5, \mu_6\mu_7) \\
&\quad + (\mu_1 - \mu_3 + \mu_5, \mu_2 + \mu_3 - \mu_4 - \mu_5 + \mu_7, \mu_6 - \mu_7)) \\
&\cong \mathbb{Z}[x, y, z, w] / (x^2, y(x-z), y(y-w), z(y+z-w), xz, w^2).
\end{aligned}$$

By an elementary computation using these presentations together with Lemmas 3.9 and 3.10, we obtain the following table; so $H^*(Z_i)$ for $i = 1, 2, 3$ are not isomorphic to each other.

Ring	s.v.e.	Maximal basis number
$H^*(Z_1)$	infinitely many	3
$H^*(Z_2)$	$2x-w, 2y-w, w$	1
$H^*(Z_3)$	$x, 2y-w, w$	2

Now we treat the case where $d \geq 5$. By Theorem 3.5, one can see that the vertices of P are one of the following:

$$\text{the vertices in } (Z_i), \pm e_4, \pm e_5, \pm(e_4 - e_5), \dots, \pm e_{d-1}, \pm e_d, \pm(e_{d-1} - e_d) \quad (Z_i^d)$$

for $i = 1, \dots, 4$ or

$$e_1, -e_1 + e_4, e_2, \pm e_3, \pm(e_2 - e_3), \pm e_4, \pm e_5, \pm(e_4 - e_5), \dots, \pm e_{d-1}, \pm e_d, \pm(e_{d-1} - e_d). \quad (Z_5^d)$$

Note that Z_5^d appears in the case $d \geq 5$ since e_4 and the vertices $e_2, \pm e_3, \pm(e_2 - e_3)$ cannot be replaced each other by unimodular transformation. In each case, $\{\pm e_{2k}, \pm e_{2k+1}, \pm(e_{2k} - e_{2k+1})\}$ ($2 \leq k \leq \frac{d-1}{2}$) forms P_6 and the first two vertices correspond to the apices of the bipyramid. We also note that $\{e_2, \pm e_3, \pm(e_2 - e_3)\}$ in (Z_5^d) forms P_5 . Therefore, one can see from the above that

$$Z_i^d = Z_i \times (X_{P_6})^{\frac{d-3}{2}} \quad \text{for } i = 1, 2, 3, 4 \quad \text{and} \quad Z_5^d = Y_2 \times X_{P_5} \times (X_{P_6})^{\frac{d-5}{2}}, \quad (3.6)$$

where Y_2 denotes the Fano 3-fold Y_2^3 in the previous subsection. Therefore, $H^*(Z_i^d)$ for $i = 1, \dots, 5$ are the tensor product of the cohomology rings of the direct factors in Z_i^d . Since Z_3 and Z_4 are diffeomorphic as observed before, so are Z_3^d and Z_4^d . It follows from Lemmas 3.7, 3.9 and 3.10 that we obtain the following table.

Ring	Maximal basis number
$H^*(Z_1^d)$	$3 + 3 \cdot \frac{d-3}{2}$
$H^*(Z_2^d)$	$1 + 3 \cdot \frac{d-3}{2}$
$H^*(Z_3^d)$	$2 + 3 \cdot \frac{d-3}{2}$
$H^*(Z_5^d)$	$1 + 3 \cdot \frac{d-3}{2}$

Thus, we have to check $H^*(Z_2^d) \cong H^*(Z_5^d)$ or not. Suppose that there is an isomorphism $F : H^*(Z_5^d) \cong H^*(Y_2) \otimes H^*(X_{P_5}) \otimes H^*(X_{P_6})^{\otimes \frac{d-5}{2}} \rightarrow H^*(Z_2^d) \cong H^*(Z_2) \otimes H^*(X_{P_6})^{\otimes \frac{d-3}{2}}$. We note that any s.v.e. of $H^*(Z_5^d)$ belongs to one of the factors of the tensor products by Lemma 3.10. Let f be an s.v.e. of $H^*(Z_5^d)$ which belongs to the factor $H^*(Y_2)$. We consider the following set

$$S(f) := \{g \in H^2(Z_5^d) \mid g \text{ is an s.v.e. and } \{f, g\} \text{ is not a part of a } \mathbb{Z}\text{-basis of } H^2(Z_5^d)\}.$$

If $g \in S(f)$, then g must belong to $H^*(Y_2)$ because otherwise $\{f, g\}$ is a part of a \mathbb{Z} -basis of $H^2(Z_5^d)$. Therefore $S(f)$ is a finite set by (3.5) and hence so is $S(F(f))$ because F is an isomorphism. This together with Lemma 3.8 shows that $F(f)$ must be an s.v.e. of $H^*(Z_2)$. This means that F sends the set of s.v.e. of $H^*(Y_2)$ to the set of s.v.e. of $H^*(Z_2)$. However, the cardinality of the former set up to sign is 4 by (3.5) while that of the latter set up to sign is 3 (see the previous subsection). This contradicts the injectivity of F . Hence, $H^*(Z_2^d)$ and $H^*(Z_5^d)$ are not isomorphic.

3.4. The case where $V(P) = 3d - 2$ with d even ≥ 4 . Theorem 3.5 says that there are three types of P 's, i.e. (1), (2) and (3) in the theorem, but they have different face numbers (Remark 3.6). Therefore, the toric Fano d -folds in these different types can be distinguished by their cohomology rings. Since there are only one smooth Fano d -polytope in (2) and (3), it suffices to investigate the case (1).

We first treat the case where $d = 4$. One can see that the vertices of P in Theorem.3.5(1) are one of the following:

$$e_1, \quad -e_1, \quad e_2, \quad -e_2, \quad \pm e_3, \pm e_4, \pm(e_3 - e_4), \quad (W_1)$$

$$e_1, \quad -e_1 + e_2, \quad e_2, \quad -e_2, \quad \pm e_3, \pm e_4, \pm(e_3 - e_4), \quad (W_2)$$

$$e_1, \quad -e_1 + e_2, \quad e_2, \quad -e_2 + e_3, \quad \pm e_3, \pm e_4, \pm(e_3 - e_4), \quad (W_3)$$

$$e_1, \quad -e_1 + e_3, \quad e_2, \quad -e_2, \quad \pm e_3, \pm e_4, \pm(e_3 - e_4), \quad (W_4)$$

$$e_1, \quad -e_1 + e_3, \quad e_2, \quad -e_2 + e_3, \quad \pm e_3, \pm e_4, \pm(e_3 - e_4), \quad (W_5)$$

$$e_1, \quad -e_1 + e_3, \quad e_2, \quad -e_2 + e_4, \quad \pm e_3, \pm e_4, \pm(e_3 - e_4), \quad (W_6)$$

$$e_1, \quad -e_1 + e_3, \quad e_2, \quad -e_2 - e_3, \quad \pm e_3, \pm e_4, \pm(e_3 - e_4), \quad (W_7)$$

$$e_1, \quad -e_1 + e_3, \quad e_2, \quad -e_2 - e_4, \quad \pm e_3, \pm e_4, \pm(e_3 - e_4). \quad (W_8)$$

In each case, $\{\pm e_3, \pm e_4, \pm(e_3 - e_4)\}$ forms the hexagon P_6 and the first two vertices and second two vertices correspond to the apices of the double bipyramid over P_6 . Note that those W_1, \dots, W_8 can be obtained by considering the pair of the first two vertices and the second two vertices, i.e. where each of two segments coming from apices intersect. We can check that those are exactly the possible cases.

We claim that W_5 is diffeomorphic to W_7 . Indeed, the vertices in W_5 are unimodularly equivalent to

$$e_1, \quad -e_1 + e_3, \quad -e_2, \quad e_2 + e_3, \quad \pm e_3, \pm e_4, \pm(e_3 - e_4)$$

through an automorphism $(x_1, x_2, x_3, x_4) \rightarrow (x_1, -x_2, x_3, x_4)$ of \mathbb{Z}^4 , and these vectors agree with the vectors in W_7 up to sign. Therefore, W_5 and W_7 are diffeomorphic by Lemma 2.3. The same argument shows that W_6 is diffeomorphic to W_8 .

We shall observe that $H^*(W_i)$ for $i = 1, \dots, 6$ are not isomorphic to each other. One can easily see

$$H^*(W_1) \cong \mathbb{Z}[x, y]/(x^2, y^2) \otimes H^*(X_{P_6}), \quad \text{and} \quad H^*(W_2) \cong \mathbb{Z}[x, y]/(x^2, y(y-x)) \otimes H^*(X_{P_6}).$$

To describe the cohomology rings of the remaining ones, we number the ten vertices

$$e_3, \quad e_4, \quad -e_3 + e_4, \quad -e_3, \quad -e_4, \quad e_3 - e_4, \quad e_1, \quad -e_1 + *, \quad e_2, \quad -e_2 + \star$$

from 1 to 10, where $*$ = e_2 or e_3 and \star = $0, \pm e_3$ or $\pm e_4$. We denote the corresponding elements in $H^2(W_i)$ for $i = 3, 4, \dots, 8$ by μ_1, \dots, μ_{10} and set

$$x = \mu_3, \quad y = \mu_4, \quad z = \mu_5, \quad w = \mu_6, \quad v = \mu_8, \quad u = \mu_{10}.$$

Then we have

$$\begin{aligned}
H^*(W_3) &\cong \mathbb{Z}[\mu_1, \mu_2, \mu_3, \mu_4, \mu_5, \mu_6, \mu_7, \mu_8, \mu_9, \mu_{10}] / \\
&\quad ((\mu_1\mu_3, \mu_1\mu_4, \mu_1\mu_5, \mu_2\mu_4, \mu_2\mu_5, \mu_2\mu_6, \mu_3\mu_5, \mu_3\mu_6, \mu_4\mu_6, \mu_7\mu_8, \mu_9\mu_{10}) \\
&\quad + (\mu_1 - \mu_3 - \mu_4 + \mu_6 + \mu_{10}, \mu_2 + \mu_3 - \mu_5 - \mu_6, \mu_7 - \mu_8, \mu_8 + \mu_9 - \mu_{10})) \\
&\cong \mathbb{Z}[x, y, z, w, v, u] / (x(x + y - u), y(x + y - u), z(y - w - u), \\
&\quad y(x - z), z(z + w), w(z + w), xz, xw, yw, v^2, u(u + v)).
\end{aligned}$$

$$\begin{aligned}
H^*(W_4) &\cong \mathbb{Z}[\mu_1, \mu_2, \mu_3, \mu_4, \mu_5, \mu_6, \mu_7, \mu_8, \mu_9, \mu_{10}] / \\
&\quad ((\mu_1\mu_3, \mu_1\mu_4, \mu_1\mu_5, \mu_2\mu_4, \mu_2\mu_5, \mu_2\mu_6, \mu_3\mu_5, \mu_3\mu_6, \mu_4\mu_6, \mu_7\mu_8, \mu_9\mu_{10}) \\
&\quad + (\mu_1 - \mu_3 - \mu_4 + \mu_6 + \mu_8, \mu_2 + \mu_3 - \mu_5 - \mu_6, \mu_7 - \mu_8, \mu_9 - \mu_{10})) \\
&\cong \mathbb{Z}[x, y, z, w, v, u] / (x(x + y - v), y(x + y - v), z(y - w - v), \\
&\quad y(x - z), z(z + w), w(z + w), xz, xw, yw, v^2, u^2).
\end{aligned}$$

$$\begin{aligned}
H^*(W_5) &\cong \mathbb{Z}[\mu_1, \mu_2, \mu_3, \mu_4, \mu_5, \mu_6, \mu_7, \mu_8, \mu_9, \mu_{10}] / \\
&\quad ((\mu_1\mu_3, \mu_1\mu_4, \mu_1\mu_5, \mu_2\mu_4, \mu_2\mu_5, \mu_2\mu_6, \mu_3\mu_5, \mu_3\mu_6, \mu_4\mu_6, \mu_7\mu_8, \mu_9\mu_{10}) \\
&\quad + (\mu_1 - \mu_3 - \mu_4 + \mu_6 + \mu_8 + \mu_{10}, \mu_2 + \mu_3 - \mu_5 - \mu_6, \mu_7 - \mu_8, \mu_9 - \mu_{10})) \\
&\cong \mathbb{Z}[x, y, z, w, v, u] / (x(x + y - v - u), y(x + y - v - u), z(y - w - v - u), \\
&\quad y(x - z), z(z + w), w(z + w), xz, xw, yw, v^2, u^2).
\end{aligned}$$

$$\begin{aligned}
H^*(W_6) &\cong \mathbb{Z}[\mu_1, \mu_2, \mu_3, \mu_4, \mu_5, \mu_6, \mu_7, \mu_8, \mu_9, \mu_{10}] / \\
&\quad ((\mu_1\mu_3, \mu_1\mu_4, \mu_1\mu_5, \mu_2\mu_4, \mu_2\mu_5, \mu_2\mu_6, \mu_3\mu_5, \mu_3\mu_6, \mu_4\mu_6, \mu_7\mu_8, \mu_9\mu_{10}) \\
&\quad + (\mu_1 - \mu_3 - \mu_4 + \mu_6 + \mu_8, \mu_2 + \mu_3 - \mu_5 - \mu_6 + \mu_{10}, \mu_7 - \mu_8, \mu_9 - \mu_{10})) \\
&\cong \mathbb{Z}[x, y, z, w, v, u] / (x(x + y - v), y(x + y - v), z(y - w - v), \\
&\quad y(x - z + u), z(z + w - u), w(z + w - u), xz, xw, yw, v^2, u^2).
\end{aligned}$$

By an elementary computation using the above presentations together with Lemma 3.7, we obtain the following table; so $H^*(W_i)$ for $i = 1, 2, 3$ are not isomorphic to each other.

Ring	s.v.e.	Maximal basis number
$H^*(W_1)$	infinitely many	5
$H^*(W_2)$	infinitely many	4
$H^*(W_3)$	$z + w, v + 2u, v$	2
$H^*(W_4)$	$2y + 2z - v, 2x + 2y - v, v, z + w, u$	3
$H^*(W_5)$	$z + w, v, u$	3
$H^*(W_6)$	$2x + 2y - v, v, 2z + 2w - u, u$	2

Now we treat the case where $d \geq 6$. The vertices of P in Theorem 3.5 (1) are either

$$\text{the vertices in } (W_i), \pm e_5, \pm e_6, \pm(e_5 - e_6), \dots, \pm e_{d-1}, \pm e_d, \pm(e_{d-1} - e_d) \quad (W_i^d)$$

for $i = 1, \dots, 8$ or

$$e_1, -e_1 + e_3, e_2, -e_2 + e_5, \pm e_3, \pm e_4, \pm(e_3 - e_4), \dots, \pm e_{d-1}, \pm e_d, \pm(e_{d-1} - e_d). \quad (W_9^d)$$

Note that W_9^d appears in the case $d \geq 6$ since e_5 and the vertices $\pm e_3, \pm e_4, \pm(e_3 - e_4)$ cannot be replaced each other by unimodular transformation. The first two vertices and second

two vertices correspond to apices of the double bipyramid and $\{\pm e_{2k-1}, \pm e_{2k}, \pm(e_{2k-1} - e_{2k})\}$ ($3 \leq k \leq \frac{d}{2}$) forms P_6 . One can see from the above that

$$W_i^d = W_i \times (X_{P_6})^{\frac{d-4}{2}} \quad \text{for } i = 1, \dots, 8 \quad \text{and} \quad W_9^d = Y_2 \times Y_2 \times (X_{P_6})^{\frac{d-6}{2}}, \quad (3.7)$$

where Y_2 is the 3-fold Y_2^3 in subsection 1.2. Since W_5 (resp. W_6) is diffeomorphic to W_7 (resp. W_8), W_5^d (resp. W_6^d) is diffeomorphic to W_7^d (resp. W_8^d). The maximal basis number of $H^*(X_{P_6})$ is 3 by Lemma 3.7 and that of $H^*(Y_2)$ is 2 by (3.5). Therefore, it follows from (3.7), Table 3.4, and Lemma 3.10 that we obtain the following table.

Ring	Maximal basis number
$H^*(W_1^d)$	$5 + 3 \cdot \frac{d-4}{2}$
$H^*(W_2^d)$	$4 + 3 \cdot \frac{d-4}{2}$
$H^*(W_3^d)$	$2 + 3 \cdot \frac{d-4}{2}$
$H^*(W_4^d)$	$3 + 3 \cdot \frac{d-4}{2}$
$H^*(W_5^d)$	$3 + 3 \cdot \frac{d-4}{2}$
$H^*(W_6^d)$	$2 + 3 \cdot \frac{d-4}{2}$
$H^*(W_9^d)$	$-2 + 3 \cdot \frac{d-4}{2}$

Thus, we have to check whether $H^*(W_3^d) \cong H^*(W_6^d)$ or not and $H^*(W_4^d) \cong H^*(W_5^d)$ or not, but the same argument as in the last part of the previous subsection shows that they are not isomorphic. Indeed, if there is an isomorphism $F : H^*(W_6^d) \rightarrow H^*(W_3^d)$, then F must send the set of s.v.e. of $H^*(W_6)$ to that of $H^*(W_3)$. However, the cardinality of the former set up to sign is 4 while that of the latter set is 3 (see Table 3.4). This contradicts the injectivity of F . Therefore, $H^*(W_3^d)$ is not isomorphic to $H^*(W_6^d)$. The same argument shows that $H^*(W_4^d)$ is not isomorphic to $H^*(W_5^d)$.

4. THE CASE OF DIMENSION 3

In the remaining sections, we use the database by Øbro mentioned in Introduction. Each smooth Fano polytope or toric Fano manifold has ID. We will denote the toric Fano manifold with ID number q by X_q .

There are 18 variety-isomorphism classes of toric Fano 3-folds, in other words, 18 unimodular equivalence classes of smooth Fano 3-polytopes. In this section, we will classify them up to diffeomorphism. It turns out that the cohomological rigidity holds for them. More precisely, there are 16 diffeomorphism classes as is shown in Table 1 below, where ID numbers whose toric Fano 3-folds are diffeomorphic are enclosed by curly braces, and five ID numbers before || are Bott manifolds. In Table 1, $V(P)$ is the number of vertices of P , P shows the unimodular equivalence classes of smooth Fano 3-polytopes, H^* shows the isomorphism classes of integer cohomology rings, and Diff shows the diffeomorphism classes.

$V(P)$	P	H^*	Diff	ID
4	1	1	1	23
5	4	4	4	7, 19, 20, 22
6	7	6	6	{11, 18}, 12, 17, 21 6, 16
7	4	3	3	8, {10, 13}, 14
8	2	2	2	9, 15
total	18	16	16	

TABLE 1. Diffeomorphism classification of toric Fano 3-folds

We shall explain how we obtain Table 1. There is only one smooth Fano 3-polytope P with $V(P) = 4$, so there is nothing to prove in this case. The case where $V(P) = 7$ or 8 is treated in Section 3. Indeed, toric Fano 3-folds X_q with $q = 8, 10, 13, 14, 9, 15$ are respectively $Z_2, Z_3, Z_4, Z_1, Y_2^3, Y_1^3$ in Section 3. Therefore, it suffices to investigate the case where $V(P) = 5$ or 6.

Convention.

- (1) The vertices of a smooth Fano 3-polytope P are shown in the database of Øbro and we number them as $1, 2, \dots$ in the order shown in the database.
- (2) The first three vertices of P are the standard basis of \mathbb{Z}^3 , so we omit them and write the vertices from 4th in Tables 2 and 4 below.
- (3) Minimal nonfaces of P are described using the numbering of the vertices of P .
- (4) \mathcal{I} denotes the ideal of the cohomology ring $H^*(X_q)$ and its minimal generators are described in the tables.
- (5) s.v.e. and c.v.e. in the tables are up to sign unless the coefficient is $\mathbb{Z}/2$.

4.1. The case where $V(P) = 5$. In this case, there are four smooth Fano 3-polytopes as shown in Table 2. They are all combinatorially equivalent to a direct sum of a 2-simplex and a 1-simplex, so the corresponding toric Fano 3-folds are generalized Bott manifolds.

ID	vertices of P from 4th	minimal nonfaces
7	$(-1, -1, 2), (0, 0, -1)$	35, 124
19	$(-1, 0, 1), (0, -1, -1)$	14, 235
20	$(-1, -1, 1), (0, 0, -1)$	35, 124
22	$(-1, 0, 0), (0, -1, -1)$	14, 235

TABLE 2. Vertices and minimal nonfaces of P with $V(P) = 5$

We denote the degree two cohomology element corresponding to 4th and 5th vertices by x and y , respectively. Then the cohomology ring of each toric Fano 3-fold with ID number in Table 2 is the quotient of a polynomial ring $\mathbb{Z}[x, y]$ by an ideal \mathcal{I} . By an elementary computation, we obtain Table 3 which shows that those four cohomology rings are not isomorphic to each other.

ID	\mathcal{I}	s.v.e.	s.v.e. $\mathbb{Z}/2$	c.v.e. $\mathbb{Z}/3$
7	$x^3, y(2x - y)$	\emptyset	(y)	
19	$x^2, y^2(x - y)$	x	(x)	(x)
20	$x^3, y(x - y)$	\emptyset	\emptyset	
22	x^2, y^3	x	(x)	(x, y)

TABLE 3. Ideals and invariants when $V(P) = 5$

4.2. The case where $V(P) = 6$. In this case, there are seven smooth Fano 3-polytopes P as shown in Table 4. The polytopes with ID numbers 11, 12, 18, 18, 21 are combinatorially equivalent to a cross-polytope, so the corresponding toric Fano 3-folds are Bott manifolds.

ID	vertices of P from 4th	minimal nonfaces
11	$(-1, 0, 1), (0, -1, 1), (0, 0, -1)$	14, 25, 36
12	$(-1, 0, 1), (0, -1, 0), (0, 1, -1)$	14, 25, 36
17	$(-1, 0, 1), (0, -1, 0), (0, 0, -1)$	14, 25, 36
18	$(-1, 0, 1), (0, -1, -1), (0, 0, -1)$	14, 25, 36
21	$(-1, 0, 0), (0, -1, 0), (0, 0, -1)$	14, 25, 36
6	$(-1, -1, 2), (0, 1, -1), (0, 0, -1)$	26, 35, 36, 124, 145
16	$(-1, 0, 1), (1, 0, -1), (-1, -1, 0)$	14, 35, 45, 126, 236

TABLE 4. Vertices and minimal nonfaces of P with $V(P) = 6$

Remark 4.1. We interchanged 5th and 6th vertices in [20] for ID numbers 12 and 18 so that the minimal nonfaces have the same numbering as others.

Through an automorphism $(x_1, x_2, x_3) \rightarrow (x_1, -x_2, x_3)$ of \mathbb{Z}^3 , the vertices of ID number 11 are unimodularly equivalent to $(1, 0, 0), (0, -1, 0), (0, 0, 1), (-1, 0, 1), (0, 1, 1), (0, 0, -1)$ and these agree with the vertices of ID number 18 up to sign, so X_{11} is diffeomorphic to X_{18} by Lemma 2.3.

We denote the degree two cohomology element corresponding to 4th, 5th and 6th vertices by x, y and z respectively. Then the cohomology ring of each toric Fano 3-fold with ID number in Table 4 is the quotient of a polynomial ring $\mathbb{Z}[x, y, z]$ by an ideal \mathcal{I} . By an elementary computation, we obtain Table 5, which shows that the cohomology rings in the tables are not isomorphic to each other:

ID	\mathcal{I}	s.v.e.	maximal basis number
11(18)	$x^2, y^2, z(x+y-z)$	x, y	2
12	$x^2, y(y-z), z(x-z)$	$x, x-2z$	1
17	$x^2, y^2, z(x-z)$	$x, y, x-2z$	2
21	x^2, y^2, z^2	x, y, z	3
6	$z(x-y), y(2x-y-z), z(x-z), x^3, x^2y$	\emptyset	0
16	$x(x+z), y^2, xy, z^3, z^2(x-y)$	y	1

TABLE 5. Ideals and invariants when $V(P) = 6$

5. THE CASE OF DIMENSION 4

There are 124 variety-isomorphism classes of toric Fano 4-folds, in other words, 124 unimodular equivalence classes of smooth Fano polytopes P of dimension 4. In this section we will classify them up to diffeomorphism. It turns out that the cohomological rigidity holds for them except for X_{50} and X_{57} . The X_{50} and X_{57} have isomorphic cohomology rings and their total Pontryagin classes are preserved under an isomorphism between their cohomology rings but we do not know whether they are diffeomorphic or not.

In Table 6 below, ID numbers whose toric Fano 4-folds are diffeomorphic are enclosed by curly braces as before. Thirteen toric Fano 4-folds with the ID numbers in the upper two lines in the row of $(V(P), F(P)) = (8, 16)$ are Bott manifolds and toric Fano 4-folds for $(V(P), F(P)) = (6, 8), (6, 9), (7, 12)$ are generalized Bott manifolds, where $V(P)$ denotes the number of vertices of P as before and $F(P)$ denotes the number of facets of P .

$V(P)$	$F(P)$	P	H^*	Diff	ID
5	5	1	1	1	147
6	8	5	5	5	25, 138, 139, 144, 145
6	9	4	3	3	44, {70, 141}, 146
7	11	3	3	3	24, 127, 128
7	12	19	16	16	{30, 43}, 31, 35, 42, 49, 66, {68, 134}, 109 117, {129, 136}, 132, 133, 135, 140, 143, 197
7	13	6	6	6	40, 41, 60, 64, 69, 137
8	15	10	7	7	26, {28, 32}, 45, 48, {67, 118}, {123, 125}, 124
8	16	28	23	23	{74, 96}, 75, {83, 108}, {95, 131} 105, 106, 112, 114, 130, 142 {29, 39}, 33, 34, 37, 38, 47, 59 93, 94, 104, {111, 116}, 115, 126
8	17	7	6	6 or 7	36, (50, 57), 58, 61, 65, 110
8	18	2	2	2	53, 55
9	18	4	4	4	27, 46, 119, 122
9	20	17	10	10	71, {73, 76, 92}, {77, 88}, 79, {81, 103} {82, 91, 107}, 84, {90, 113}, 102, 120
9	21	4	4	4	51, 52, 56, 89
9	23	1	1	1	62
9	24	1	1	1	54
10	24	8	6	6	{72, 87}, {78, 86}, 80, 85, 101, 121
10	25	1	1	1	98
10	30	1	1	1	63
11	30	1	1	1	99
12	36	1	1	1	100
total		124	102	102 or 103	

TABLE 6. Diffeomorphism classification of toric Fano 4-folds

Our approach to obtain the table above is the same as the case of dimension 3 but the analysis of the cohomology rings in dimension 4 becomes much more complicated and there are many more cases to investigate.

When $V(P) = 5, 11, 12$ or $(V(P), F(P)) = (9, 23), (9, 24), (10, 25), (10, 30)$, there is only one smooth Fano 4-polytope; so there is nothing to prove in these cases. Moreover, the case $(V(P), F(P)) = (10, 24)$ is treated in Section 3. Indeed, toric Fano 4-folds with ID numbers 72, 78, 80, 85, 86, 87, 101, 121 are respectively $W_5, W_6, W_3, W_4, W_8, W_7, W_2, W_1$ in Subsection 3.4. Therefore, it suffices to investigate the remaining cases. We shall carry out this task one by one in this section.

Convention.

- (1) The vertices of a smooth Fano 4-polytope P are shown in the database by \emptyset bro and we number them as 1, 2, ... in the order shown in the database.
- (2) The first four vertices of P are the standard basis of \mathbb{Z}^4 , so we omit them and write the vertices from 5th in the tables below.
- (3) Minimal nonfaces of P are described using the numbering of the vertices of P .
- (4) \mathcal{I} denotes the ideal of the cohomology ring $H^*(X_q)$ and its minimal generators are described in the tables.
- (5) s.v.e., c.v.e. and 4-v.e. in the tables are up to sign unless the coefficient is $\mathbb{Z}/2$.
- (6) $X_p \cong X_q$ means that X_p is diffeomorphic to X_q .
- (7) $H^*(X_p) \not\cong H^*(X_q)$ means that the cohomology rings are not isomorphic (as graded rings).
- (8) The degree two cohomology elements corresponding to 5th, 6th, 7th, 8th, 9th vertices are respectively denoted by x, y, z, u, v .

5.1. **The case where $V(P) = 6$.** We take two cases according to the values of $F(P)$.

5.1.1. $(V(P), F(P)) = (6, 8)$. In this case, there are five smooth Fano 4-polytopes and they are all combinatorially equivalent to a direct sum of a 3-simplex and a 1-simplex, so the corresponding toric Fano 4-folds are generalized Bott manifolds. Using the data in Table 7, we obtain Table 8 which shows that the five cohomology rings are not isomorphic to each other.

ID	vertices of P from 5th	minimal nonfaces
25	$(-1, -1, -1, 3), (0, 0, 0, -1)$	1235, 46
138	$(-1, 0, 0, 1), (0, -1, -1, -1)$	2346, 15
139	$(-1, -1, -1, 2), (0, 0, 0, -1)$	1235, 46
144	$(-1, 0, 0, 0), (1, -1, -1, -1)$	2346, 15
145	$(-1, 0, 0, 0), (0, -1, -1, -1)$	2346, 15

TABLE 7. Vertices and minimal nonfaces of P with $(V(P), F(P)) = (6, 8)$

ID	\mathcal{I}	s.v.e.	s.v.e. $\mathbb{Z}/2$	4-v.e. $\mathbb{Z}/2$	c.v.e. $\mathbb{Z}/3$
25	$x^4, y(y-3x)$	\emptyset	\emptyset	x	y
138	$x^4, y(y-x)$	\emptyset	\emptyset	x	\emptyset
139	$x^4, y(y-2x)$	\emptyset	y		
144	$x^3(x-y), y^2$	y	y	y	
145	x^4, y^2	y	y	(x, y)	

TABLE 8. Ideals and invariants when $(V(P), F(P)) = (6, 8)$

5.1.2. $(V(P), F(P)) = (6, 9)$. In this case, there are four smooth Fano 4-polytopes and they are all combinatorially equivalent to a direct sum of two 2-simplices, so the corresponding toric Fano 4-folds are generalized Bott manifolds.

ID	vertices of P from 5th	minimal nonfaces
44	$(-1, -1, 0, 2), (0, 0, -1, -1)$	125, 346
70	$(-1, -1, 1, 1), (0, 0, -1, -1)$	125, 346
141	$(-1, -1, 0, 1), (0, 0, -1, -1)$	125, 346
146	$(-1, -1, 0, 0), (0, 0, -1, -1)$	125, 346

TABLE 9. Vertices and minimal nonfaces of P with $(V(P), F(P)) = (6, 9)$

We see that

$$X_{70} \cong X_{141}, \quad (5.1)$$

see Subsection 5.5. On the other hand, using the data in Table 9, we obtain Table 10 which shows that the three cohomology rings are not isomorphic to each other.

ID	\mathcal{I}	c.v.e.	c.v.e. $\mathbb{Z}/2$
44	$x^3, y^2(2x-y)$	x	x, y
70(141)	$x^3, y(x-y)^2$	x	x
146	x^3, y^3	x, y	

TABLE 10. Ideals and invariants when $(V(P), F(P)) = (6, 9)$

5.2. **The case where $V(P) = 7$.**

5.2.1. $(V(P), F(P)) = (7, 11)$. In this case, there are three smooth Fano 4-polytopes and they are all combinatorially equivalent. Using the data in Table 11, we obtain Table 12 which shows that the three cohomology rings are not isomorphic to each other.

ID	vertices of P from 5th	minimal nonfaces
24	$(-1, -1, -1, 3), (0, 0, 1, -1), (0, 0, 0, -1)$	1235, 1256, 37, 46, 47
127	$(-1, 0, 0, 1), (1, 0, 0, -1), (-1, -1, -1, 0)$	1237, 2347, 15, 46, 56
128	$(-1, 0, 0, 1), (-1, 0, 0, 0), (2, -1, -1, -1)$	2347, 2357, 15, 16, 46

TABLE 11. Vertices and minimal nonfaces of P with $(V(P), F(P)) = (7, 11)$

ID	\mathcal{I}	s.v.e.	c.v.e. $\mathbb{Z}/3$
24	$z(x-y), y(y+z-3x), z(z-2x), x^4, x^3y$	\emptyset	$y+z$
127	$x(x+z), y^2, xy, z^4, z^3(y-x)$	y	
128	$x(x+y-2z), y(y-z), y(z-x), z^4, xz^3$	\emptyset	\emptyset

TABLE 12. Ideals and invariants when $(V(P), F(P)) = (7, 11)$

5.2.2. $(V(P), F(P)) = (7, 12)$. In this case, there are 19 smooth Fano 4-polytopes and they are all combinatorially equivalent to a direct sum of 2-simplex and two 1-simplices, so the corresponding toric Fano 4-folds are generalized Bott manifolds.

ID	vertices of P from 5th	minimal nonfaces
30	$(-1, -1, 0, 2), (0, 0, -1, 1), (0, 0, 0, -1)$	125, 36, 47
31	$(-1, -1, 0, 2), (0, 0, 1, -1), (0, 0, -1, 0)$	125, 37, 46
35	$(-1, -1, 0, 2), (0, 1, -1, 0), (0, 0, 0, -1)$	125, 36, 47
42	$(-1, -1, 0, 2), (0, 0, -1, 0), (0, 0, 0, -1)$	125, 36, 47
43	$(-1, -1, 0, 2), (0, 0, 0, -1), (0, 0, -1, -1)$	125, 37, 46
49	$(-1, -1, 1, 1), (0, 0, -1, 1), (0, 0, 0, -1)$	125, 36, 47
66	$(-1, -1, 1, 1), (0, 0, -1, 0), (0, 0, 0, -1)$	125, 36, 47
68	$(-1, -1, 1, 1), (0, 0, -1, 0), (0, 0, -1, -1)$	125, 36, 47
97	$(-1, 0, 0, 1), (0, -1, 0, 1), (0, 0, -1, -1)$	347, 15, 26
109	$(-1, 0, 0, 1), (0, -1, 1, 0), (0, 0, -1, -1)$	347, 15, 26
117	$(-1, 0, 0, 1), (0, 0, 1, -1), (0, -1, -1, 0)$	237, 15, 46
129	$(-1, 0, 0, 1), (0, -1, -1, 1), (0, 0, 0, -1)$	236, 15, 47
132	$(-1, 0, 0, 1), (0, -1, 0, 0), (0, 1, -1, -1)$	347, 15, 26
133	$(-1, 0, 0, 1), (0, -1, 0, 0), (0, 0, -1, -1)$	347, 15, 26
134	$(-1, 0, 0, 1), (0, 0, 0, -1), (1, -1, -1, 0)$	237, 15, 46
135	$(-1, 0, 0, 1), (0, 0, 0, -1), (0, -1, -1, 0)$	237, 15, 46
136	$(-1, 0, 0, 1), (0, 0, 0, -1), (0, -1, -1, -1)$	237, 15, 46
140	$(-1, -1, 0, 1), (0, 0, -1, 0), (0, 0, 0, -1)$	125, 36, 47
143	$(-1, 0, 0, 0), (0, -1, 0, 0), (0, 0, -1, -1)$	347, 15, 26

TABLE 13. Vertices and minimal nonfaces of P with $(V(P), F(P)) = (7, 12)$

We see that

$$X_{30} \cong X_{43}, \quad X_{68} \cong X_{134}, \quad X_{129} \cong X_{136}, \quad (5.2)$$

see Subsection 5.5. Using the data in Table 13, we obtain the following table.

ID	\mathcal{I}	s.v.e.	s.v.e. $\mathbb{Z}/2$	4-v.e. $\mathbb{Z}/2$	c.v.e. $\mathbb{Z}/3$	c.v.e. $\mathbb{Z}/2$
30 (43)	$x^3, y^2, z(2x + y - z)$	y	y	all	(x, y)	$x, y, z,$ $y + z$
31	$x^3, y(2x - y), z(y - z)$	\emptyset	y			
35	$x^2(x - y), y^2, z(2x - z)$	y	(y, z)	all	y	
42	$x^3, y^2, z(2x - z)$	y	(y, z)	all	(x, y)	
49	$x^3, y(x - y), z(x + y - z)$	\emptyset	\emptyset	(x, y)	x	x
66	$x^3, y(x - y), z(x - z)$	\emptyset	\emptyset	all		
68 (134)	$x^3, y(x - y - z), z(x - z)$	\emptyset	\emptyset	(x, z)	x	x
97	$x^2, y^2, z^2(x + y - z)$	x, y	(x, y)	all	(x, y)	$x, y, x + y,$ $x + y + z$
109	$x^2, y^2, z(x - z)(y - z)$	x, y	(x, y)	(x, y)		
117	$x^2, y(x - y), z^2(y - z)$	$x, x - 2y$	x	(x, y)		
129 (136)	$x^2, y^3, z(x + y - z)$	x	x	(x, y)	(x, y)	
132	$x^2, y(y - z), z^2(x - z)$	x	x	(x, z)	x	
133	$x^2, y^2, z^2(x - z)$	x, y	(x, y)	all	(x, y)	$x, y, x + y$ $x + z$
135	$x^2, y(x - y), z^3$	$x, x - 2y$	x	all		
140	$x^3, y^2, z(x - z)$	y	y	all	(x, y)	x, y
143	x^2, y^2, z^3	x, y	(x, y)	all	all	

TABLE 14. Ideals and invariants when $(V(P), F(P)) = (7, 12)$

Table 14 shows that the cohomology rings in the table are distinguished by the invariants in the table except for two pairs: 49 and 68(134), 97 and 133. We shall prove that their cohomology rings are not isomorphic to each other.

The c.v.e. of $H^*(X_{49})$ and $H^*(X_{68})$ are both x up to sign. Therefore, if there is an isomorphism $H^*(X_{49}) \rightarrow H^*(X_{68})$, then it induces an isomorphism $H^*(X_{49})/(x^2) \rightarrow H^*(X_{68})/(x^2)$. However, Table 15 shows that this does not occur, so $H^*(X_{49}) \not\cong H^*(X_{68})$.

ID	$\mathcal{I} + (x^2)$	c.v.e. $\mathbb{Z}/3$
49	$x^2, y(x - y), z(x + y - z)$	all
68(134)	$x^2, y(x - y - z), z(x - z)$	(x, z)

TABLE 15. Distinguishment between $H^*(X_{49})$ and $H^*(X_{68})$

The s.v.e. of $H^*(X_{97})$ and $H^*(X_{133})$ are both x, y up to sign and the transposition of x and y induces an automorphism of $H^*(X_{97})$ since the ideal \mathcal{I} of $H^*(X_{97})$ is invariant under the transposition. Therefore, if there is an isomorphism $F: H^*(X_{97}) \rightarrow H^*(X_{133})$, then we may assume that $F(x) = \pm x$, so that F induces an isomorphism $H^*(X_{97})/(x) \rightarrow H^*(X_{133})/(x)$. However, Table 16 shows that this does not occur, so $H^*(X_{97}) \not\cong H^*(X_{133})$.

ID	$\mathcal{I} + (x)$	c.v.e. $\mathbb{Z}/3$
97	$y^2, z^2(y - z)$	y
133	y^2, z^3	all

TABLE 16. Distinguishment between $H^*(X_{97})$ and $H^*(X_{133})$

5.2.3. $(V(P), F(P)) = (7, 13)$. In this case, there are six smooth Fano 4-polytopes. We shall prove that these six cohomology rings are not isomorphic to each other.

ID	vertices of P from 5th	minimal nonfaces
40	$(-1, -1, 0, 2), (0, 1, 0, -1), (0, 0, -1, -1)$	125, 156, 237, 347, 46
41	$(-1, -1, 0, 2), (1, 1, -1, -1), (0, 0, 0, -1)$	125, 127, 346, 356, 47
60	$(-1, -1, 1, 1), (0, 1, -1, 0), (0, 0, -1, -1)$	125, 156, 247, 347, 36
64	$(-1, -1, 1, 1), (1, 1, -1, -1), (-1, -1, 0, 0)$	125, 127, 346, 347, 56
69	$(-1, -1, 1, 1), (0, 1, -1, -1), (0, 0, -1, -1)$	125, 156, 346, 347, 27
137	$(-1, 0, 0, 1), (1, -1, 0, -1), (-1, 0, -1, 0)$	137, 246, 256, 347, 15

TABLE 17. Vertices and minimal nonfaces of P with $(V(P), F(P)) = (7, 13)$

Using the data in Table 17, we obtain the following table.

ID	\mathcal{I}	c.v.e.	c.v.e. $\mathbb{Z}/2$
40	$x^3, x^2y, z^2(x-y), z^2(x-z), y(2x-y-z)$	x	$x, y+z$
41	$x(x-y)^2, y(x-y)^2, y^2(y+z), xy^2, z(2x-y-z)$	\emptyset	
60	$x^3, x^2y, z(x-y)(x-z), z^2(x-z), y(x-y-z)$	x	x
64	$x(x+z)^2, z(x-y+z)^2, y^3, z(x-y)^2, xy$	y	y, z
69	$x^3, x^2y, y(x-y-z)^2, z^3, z(x-y)$	x, z	x, z
137	$z^3, y^3, xy^2, z^2(x-y), x(x-y+z)$	y, z	y, z

TABLE 18. Ideals and invariants when $(V(P), F(P)) = (7, 13)$

Table 18 shows that the six cohomology rings can be distinguished by the invariants in the table except for two pairs: 40 and 64, 69 and 137. We shall prove that their cohomology rings are not isomorphic to each other.

The c.v.e. of $H^*(X_{40})$ and $H^*(X_{64})$ are respectively x and y up to sign. Therefore, if there is an isomorphism $H^*(X_{40}) \rightarrow H^*(X_{64})$, then it induces an isomorphism $H^*(X_{40})/(x) \rightarrow H^*(X_{64})/(y)$. However, this does not occur because

$$\begin{aligned} H^*(X_{40})/(x) &= \mathbb{Z}[y, z]/(yz^2, z^3, y(y+z)), \\ H^*(X_{64})/(y) &= \mathbb{Z}[x, z]/(x(x+z)^2, z(x+z)^2, x^2z), \end{aligned}$$

and the degree sequences of these ideals are different. Therefore $H^*(X_{40}) \not\cong H^*(X_{64})$.

Similarly, the c.v.e. of $H^*(X_{69})$ and $H^*(X_{137})$ are respectively x, z and y, z up to sign. Therefore, if there is an isomorphism $H^*(X_{69}) \rightarrow H^*(X_{137})$, then it induces an isomorphism $H^*(X_{69})/(x, z) \rightarrow H^*(X_{137})/(y, z)$. However, this does not occur because

$$H^*(X_{69})/(x, z) = \mathbb{Z}[y]/(y^3), \quad H^*(X_{137})/(y, z) = \mathbb{Z}[x]/(x^2),$$

and these quotient rings are not isomorphic. Therefore $H^*(X_{69}) \not\cong H^*(X_{137})$.

5.3. The case where $V(P) = 8$.

5.3.1. $(V(P), F(P)) = (8, 15)$. In this case, there are ten smooth Fano 4-polytopes and they are all combinatorially equivalent to a direct sum of a 5-gon and a 2-simplex.

ID	vertices of P from 5th	minimal nonfaces
26	$(-1, -1, 0, 2), (0, 0, -1, 1), (0, 0, 1, -1), (0, 0, -1, 0)$	125, 36, 38, 47, 48, 67
28	$(-1, -1, 0, 2), (0, 0, -1, 1), (0, 0, 1, -1), (0, 0, 0, -1)$	125, 36, 38, 47, 48, 67
32	$(-1, -1, 0, 2), (0, 0, 1, -1), (0, 0, -1, 0), (0, 0, 0, -1)$	125, 37, 38, 46, 48, 67
45	$(-1, -1, 1, 1), (0, 0, -1, 1), (0, 0, 1, -1), (0, 0, -1, 0)$	125, 36, 38, 47, 48, 67
48	$(-1, -1, 1, 1), (0, 0, -1, 1), (0, 0, -1, 0), (0, 0, 0, -1)$	125, 36, 37, 47, 48, 68
67	$(-1, -1, 1, 1), (0, 0, -1, 0), (0, 0, 0, -1), (0, 0, -1, -1)$	125, 36, 38, 47, 48, 67
118	$(-1, 0, 0, 1), (1, 0, 0, -1), (-1, 0, 0, 0), (0, -1, -1, 1)$	238, 15, 17, 46, 47, 56
123	$(-1, 0, 0, 1), (1, 0, 0, -1), (-1, 0, 0, 0), (1, -1, -1, 0)$	238, 15, 17, 46, 47, 56
124	$(-1, 0, 0, 1), (1, 0, 0, -1), (-1, 0, 0, 0), (0, -1, -1, 0)$	238, 15, 17, 46, 47, 56
125	$(-1, 0, 0, 1), (1, 0, 0, -1), (-1, 0, 0, 0), (1, -1, -1, -1)$	238, 15, 17, 46, 47, 56

TABLE 19. Vertices and minimal nonfaces of P with $(V(P), F(P)) = (8, 15)$

We see that

$$X_{28} \cong X_{32}, \quad X_{67} \cong X_{118}, \quad X_{123} \cong X_{125}, \quad (5.3)$$

see Subsection 5.5. We obtain the following table from Table 19.

ID	\mathcal{I}	s.v.e.	s.v.e. $\mathbb{Z}/2$	c.v.e.
26	$x^3, y(y+u), u(y-z+u),$ $z(2x-z), u(2x-u), yz$	\emptyset	(z, u)	
28 (32)	$x^3, y^2, u(y-z),$ $z(2x-z-u), u(2x-u), yz$	y	(y, u)	
45	$x^3, y(x-y-u), u(2x-u),$ $x(x-z), u(x+y-z), yz$	\emptyset	u	
48	$x^3, y(x-y-z), z(x-y-z),$ $z(x+y-u), u(x-u), yu$	\emptyset	\emptyset	x
67 (118)	$x^3, y(x-y-u), u(x-y-u),$ $z(x-z-u), u(y-z), yz$	\emptyset	\emptyset	x
123 (125)	$u^3, x(x+z-u), z(z-u),$ $y^2, z(x-y), xy$	y	y	
124	$u^3, x(x+z), z^2, y^2, z(x-y), xy$	∞		

TABLE 20. Ideals and invariants when $(V(P), F(P)) = (8, 15)$

Table 20 shows that the seven cohomology rings in the table can be distinguished by the invariants in the table except for 48 and 67(118). The c.v.e. of $H^*(X_{48})$ and $H^*(X_{67})$ are both x up to sign. Therefore, if there is an isomorphism $H^*(X_{48}) \rightarrow H^*(X_{67})$, then it induces an isomorphism $H^*(X_{48})/(x^2) \rightarrow H^*(X_{67})/(x^2)$. However, Table 21 shows that this does not occur, so $H^*(X_{48}) \not\cong H^*(X_{67})$.

ID	$\mathcal{I} + (x^2)$	c.v.e. $\mathbb{Z}/3$
48	$x^2, y(x-y-z), z(x-y-z), z(x+y-u), u(x-u), yu$	(x, y, z, u)
67(118)	$x^2, (y(x-y-u), u(x-y-u), z(x-z-u), u(x-z-u), yz$	$(x, y+u, z+u)$

TABLE 21. Distinguishment between $H^*(X_{48})$ and $H^*(X_{67})$

5.3.2. $(V(P), F(P)) = (8, 16)$. In this case, there are 28 smooth Fano 4-polytopes and there are two combinatorial types among them. Indeed, 13 polytopes among them have the same combinatorial type as a cross-polytope as shown in Table 22. The corresponding toric Fano 4-folds are Bott manifolds. The cohomology rings associated to these two different combinatorial types are not isomorphic to each other because the degree sequences of their

ideals are different, see Tables 23 and 25. More generally, it is known that a toric manifold which has the same cohomology ring as a Bott manifold is indeed a Bott manifold ([18]).

ID	vertices of P from 5th	minimal nonfaces
74	$(-1, 0, 0, 1), (0, -1, 0, 1), (0, 0, -1, 1), (0, 0, 0, -1)$	15, 26, 37, 48
75	$(-1, 0, 0, 1), (0, -1, 0, 1), (0, 0, 1, -1), (0, 0, -1, 0)$	15, 26, 38, 47
83	$(-1, 0, 0, 1), (0, -1, 0, 1), (0, 1, -1, 0), (0, 0, 0, -1)$	15, 26, 37, 48
95	$(-1, 0, 0, 1), (0, -1, 0, 1), (0, 0, -1, 0), (0, 0, 0, -1)$	15, 26, 37, 48
96	$(-1, 0, 0, 1), (0, -1, 0, 1), (0, 0, 0, -1), (0, 0, -1, -1)$	15, 26, 38, 47
105	$(-1, 0, 0, 1), (0, -1, 1, 0), (0, 1, 0, -1), (0, 0, -1, 0)$	15, 26, 38, 47
106	$(-1, 0, 0, 1), (0, -1, 1, 0), (0, 0, -1, 0), (0, 0, 0, -1)$	15, 26, 37, 48
108	$(-1, 0, 0, 1), (0, -1, 1, 0), (0, 0, -1, 0), (0, 0, -1, -1)$	15, 26, 37, 48
112	$(-1, 0, 0, 1), (0, 0, 1, -1), (0, -1, 0, 0), (0, 0, -1, 0)$	15, 27, 38, 46
114	$(-1, 0, 0, 1), (0, 0, 1, -1), (0, -1, 1, -1), (0, 0, -1, 0)$	15, 27, 38, 46
130	$(-1, 0, 0, 1), (0, -1, 0, 0), (0, 0, -1, 0), (0, 0, 0, -1)$	15, 26, 37, 48
131	$(-1, 0, 0, 1), (0, -1, 0, 0), (0, 0, 0, -1), (0, 0, -1, -1)$	15, 26, 38, 47
142	$(-1, 0, 0, 0), (0, -1, 0, 0), (0, 0, -1, 0), (0, 0, 0, -1)$	15, 26, 37, 48

TABLE 22. Vertices and minimal nonfaces of P with $(V(P), F(P)) = (8, 16)$, cross-polytope

One can see that

$$X_{74} \cong X_{96}, \quad X_{83} \cong X_{108}, \quad X_{95} \cong X_{131}, \quad (5.4)$$

see Subsection 5.5. Using the data in Table 22, we obtain the following table.

ID	\mathcal{I}	s.v.e.	4-v.e. $\mathbb{Z}/2$
74(96)	$x^2, y^2, z^2, u(x+y+z-u)$	x, y, z	all
75	$x^2, y^2, u(z-u), z(x+y-z)$	x, y	all
83(108)	$x^2, y(y-z), z^2, u(x+y-u)$	$x, z, z-2y$	(x, y, z)
95(131)	$x^2, y^2, z^2, u(x+y-u)$	x, y, z	all
105	$x^2, y(y-z), u(y-u), z(x-z)$	$x, x-2z$	(x, y, z)
106	$x^2, y^2, z(y-z), u(x-u)$	$x, y, x-2u, y-2z$	
112	$x^2, z^2, u(y-u), y(x-y)$	$x, z, x-2y$	all
114	$x^2, z^2, u(y+z-u), y(x-y-z)$	x, z	(x, y, z)
130	$x^2, y^2, z^2, u(x-u)$	$x, y, z, x-2u$	
142	x^2, y^2, z^2, u^2	x, y, z, u	

TABLE 23. Ideals and invariants when $(V(P), F(P)) = (8, 16)$, cross-polytope

Table 23 shows that the cohomology rings in the table can be distinguished by the invariants in the table except for 74(96) and 95.

Suppose that there is an isomorphism $F : H^*(X_{74}) \rightarrow H^*(X_{95})$. Since the s.v.e. of $H^*(X_{74})$ and $H^*(X_{95})$ are x, y, z and any permutation of x, y, z is an automorphism of $H^*(X_{74})$, we may assume that $F(x) = \pm x, F(y) = \pm y, F(z) = \pm z$. Therefore, F induces an isomorphism $H^*(X_{74})/(x, y) \cong H^*(X_{95})/(x, y)$. However, this does not occur because

$$H^*(X_{74})/(x, y) = \mathbb{Z}[z, u]/(z^2, u(u-z)), \quad H^*(X_{95}) = \mathbb{Z}[z, u]/(z^2, u^2)$$

and these two rings are not isomorphic (e.g. their s.v.e. are different). Therefore, $H^*(X_{74}) \not\cong H^*(X_{95})$.

Next, we shall treat the other case where P is not combinatorially equivalent to a cross-polytope. There are 15 smooth Fano 4-polytopes in this case as shown in Table 24.

ID	vertices of P from 5th	minimal nonfaces
29	$(-1, -1, 0, 2), (0, 0, -1, 1), (0, 1, 0, -1), (0, 0, 0, -1)$	125, 157, 28, 36, 47, 48
33	$(-1, -1, 0, 2), (0, 1, -1, 0), (0, 1, 0, -1), (0, 0, 0, -1)$	125, 157, 28, 36, 47, 48
34	$(-1, -1, 0, 2), (0, 1, -1, 0), (1, 0, 0, -1), (0, 0, 0, -1)$	125, 257, 18, 36, 47, 48
37	$(-1, -1, 0, 2), (0, 1, 0, -1), (0, 0, -1, 0), (0, 0, 0, -1)$	125, 156, 28, 37, 46, 48
38	$(-1, -1, 0, 2), (0, 1, 0, -1), (0, 0, 0, -1), (0, 1, -1, -1)$	125, 156, 27, 38, 46, 47
39	$(-1, -1, 0, 2), (0, 1, 0, -1), (0, 0, 0, -1), (0, 0, -1, -1)$	125, 156, 27, 38, 46, 47
47	$(-1, -1, 1, 1), (0, 0, -1, 1), (0, 1, 0, -1), (0, 0, 0, -1)$	125, 157, 28, 36, 47, 48
59	$(-1, -1, 1, 1), (0, 1, -1, 0), (0, 0, -1, 0), (0, 0, 0, -1)$	125, 156, 27, 36, 37, 48
93	$(-1, 0, 0, 1), (0, -1, 0, 1), (0, 1, 0, -1), (0, -1, -1, 0)$	238, 348, 15, 26, 47, 67
94	$(-1, 0, 0, 1), (0, -1, 0, 1), (0, 1, 0, -1), (0, 0, -1, -1)$	238, 348, 15, 26, 47, 67
104	$(-1, 0, 0, 1), (0, -1, 1, 0), (0, 1, -1, 0), (0, -1, 0, -1)$	248, 348, 15, 26, 37, 67
111	$(-1, 0, 0, 1), (0, 0, 1, -1), (0, -1, -1, 1), (0, 0, 0, -1)$	237, 267, 15, 38, 46, 48
115	$(-1, 0, 0, 1), (0, 0, 1, -1), (0, 0, 0, -1), (1, -1, -1, 0)$	238, 268, 15, 37, 46, 47
116	$(-1, 0, 0, 1), (0, 0, 1, -1), (0, 0, 0, -1), (0, -1, -1, 0)$	238, 268, 15, 37, 46, 47
126	$(-1, 0, 0, 1), (1, 0, 0, -1), (0, -1, 0, 0), (-1, 0, -1, 0)$	138, 348, 15, 27, 46, 56

TABLE 24. Vertices and minimal nonfaces of P with $(V(P), F(P)) = (8, 16)$, non-cross-polytope

We see that

$$X_{29} \cong X_{39}, \quad X_{111} \cong X_{116}, \quad (5.5)$$

see Subsection 5.5. Using the data in Table 24, we obtain the following table.

ID	\mathcal{I}	s.v.e.	s.v.e. $\mathbb{Z}/2$	4-v.e. $\mathbb{Z}/2$	c.v.e. $\mathbb{Z}/3$
29 (39)	$x^3, x^2z, u(x-z), y^2,$ $z(2x+y-z-u), u(x+y-u)$	y	y	$(x, y, z+u)$	(x, y)
33	$x^2(x-y), x^2z, u(x-y-z), y^2,$ $z(2x-z-u), u(2x-z-u)$	y	$(y, z+u)$	all	y
34	$x^2(x-y), xz(x-y), u(x-z),$ $y^2, z(2x-z-u), u(x-u)$	y	$(y, z+u)$	$(x, y, z+u)$	
37	$x^3, x^2y, u(x-y), z^2,$ $y(2x-y-u), u(x-u)$	z	$(y+u, z)$	all	(x, y, z)
38	$x^2(x-u), x^2y, z(x-y-u), u^2,$ $y(2x-y-z-u), z(x-z)$	u	u	$(x, y+z, u)$	u
47	$x^3, x^2z, u(x-z), y(x-y),$ $z(x+y-z-u), u(y-u)$	\emptyset	\emptyset	(x, y, z)	x
59	$x^3, x^2y, z(x-y), y(x-y-z),$ $z^2, u(x-u)$	z	z	all	
93	$u^2(y-z+u), u^2(x-u),$ $x^2, y(y+u), z(x-z), yz$	$x, x-2z$	x	(x, z, u)	(x, z)
94	$u^2(y-z), u^2(x-u), x^2,$ $y^2, z(x-z-u), yz$	x, y	(x, y)	all	(x, y)
104	$u^2(x-u), u(y-z)(x-u),$ $x^2, y(y+u), z^2, yz$	x, z	(x, z)	(x, z, u)	
111 (116)	$z^3, yz^2, x^2, u(y-z),$ $y(x-y+z-u), u(x-u)$	$x, x-2u$	x	(x, z, u)	(x, z, u)
115	$u^3, yu^2, x(x-u), z(y-u),$ $y(x-y-z), z(x-y-z)$	\emptyset	\emptyset	(x, y, u)	u
126	$u^3, u^2(x-y), x(x+u),$ z^2, y^2, xy	y, z	(y, z)	all	(y, z, u)

TABLE 25. Ideals and invariants when $(V(P), F(P)) = (8, 16)$, non-cross-polytope

Table 25 shows that the cohomology rings in the table are distinguished by the invariants in the table except for 47 and 115. The c.v.e. of $H^*(X_{47})$ and $H^*(X_{115})$ are respectively

x and u up to sign. Therefore, if there is an isomorphism $H^*(X_{47}) \rightarrow H^*(X_{115})$, then it induces an isomorphism $H^*(X_{47})/(x^2) \rightarrow H^*(X_{115})/(u^2)$. However, Table 26 shows that this does not occur, so $H^*(X_{47}) \not\cong H^*(X_{115})$.

ID	$\mathcal{I} + ((\text{c.v.e.})^2)$	c.v.e. $\mathbb{Z}/3$
47	$x^2, u(x-z), y(x-y), z(x+y-z-u), u(y-u)$	$(x, y, z+u)$
115	$u^2, x(x-u), z(y-u), y(x-y-z), z(x-y-z)$	(x, u)

TABLE 26. Distinguishment between $H^*(X_{47})$ and $H^*(X_{115})$

5.3.3. $(V(P), F(P)) = (8, 17)$. There are seven smooth Fano 4-polytopes in this case and there are two combinatorial types among them: one is ID numbers 36, 65 and the other one is ID numbers 50, 57, 58, 61, 110, see Table 27. One can distinguish the cohomology rings of the former class between those of the latter class by the degree sequences of the ideals. Therefore, Table 28 shows that the seven cohomology rings can be distinguished by the degree sequences of the ideals and c.v.e. over $\mathbb{Z}/2$ except for 50 and 57.

The fans of X_{50} and X_{57} do not satisfy the condition in Lemma 2.3 up to unimodular equivalence, so they are not *weakly equivariantly* diffeomorphic with respect to the restricted $(S^1)^4$ -actions. However, their cohomology rings are isomorphic to each other. Indeed, the map

$$(x, y, z, u) \rightarrow (-x + 2u, -y + u, u, -z) \quad (5.6)$$

gives an isomorphism $H^*(X_{50}) \rightarrow H^*(X_{57})$. It does not preserve their total Chern classes (even their first Chern classes) but it does preserve their total Pontryagin classes (in fact, the first and second Pontryagin classes because of dimensional reason) which are given by

$$\begin{aligned} p(X_{50}) &= (1+x^2)^2(1+(x-y-z+u)^2)(1+(x-y)^2)(1+(x-z)^2)(1+y^2)(1+z^2)(1+u^2), \\ p(X_{57}) &= (1+x^2)^2(1+(x-y+u)^2)(1+(x-y)^2)(1+(x-z)^2)(1+y^2)(1+z^2)(1+u^2), \end{aligned}$$

see Remark 2.2.

ID	vertices of P from 5th	minimal nonfaces
36	$(-1, -1, 0, 2), (0, 1, 0, -1), (0, -1, -1, 1), (0, 0, 0, -1)$	28, 46, 48, 125, 156, 158, 237, 347, 367
50	$(-1, -1, 1, 1), (0, 1, -1, 0), (0, 1, 0, -1), (0, -1, 0, 0)$	28, 36, 47, 125, 156, 157, 348
57	$(-1, -1, 1, 1), (0, 1, -1, 0), (0, -1, 0, 0), (0, 0, 0, -1)$	27, 36, 48, 125, 156, 158, 347
58	$(-1, -1, 1, 1), (0, 1, -1, 0), (0, -1, 0, 0), (0, 1, -1, -1)$	27, 36, 48, 125, 156, 158, 347
61	$(-1, -1, 1, 1), (1, 1, -1, -1), (-1, 0, 0, 0), (0, -1, 0, 0)$	17, 28, 56, 125, 346, 347, 348
65	$(-1, -1, 1, 1), (1, 1, -1, -1), (-1, -1, 0, 0), (0, 0, -1, -1)$	56, 58, 67, 125, 127, 128, 346, 347, 348
110	$(-1, 0, 0, 1), (0, 0, 1, -1), (1, 0, -1, 0), (-1, -1, 0, 0)$	15, 37, 46, 128, 238, 248, 567

TABLE 27. $(V(P), F(P)) = (8, 17)$

ID	\mathcal{I}	c.v.e. $\mathbb{Z}/2$
36	$(x - y + z)u, (2x - y + z - u)y, (x - u)u, x^2(x + z),$ $x^2y, x^2u, (x + z)z^2, z^2(x - u), z^2y$	$x + z, (y + z, u)$
65	$xy, xu, yz, (x + z)^2x, (x + z)^2z,$ $(y - z)^2u, (x - y - u)^2y, (x - u)^2z, (y + u)^2u$	$z + u, y + u, x + z$
50	$(x - y - z + u)u, (x - y)y, (x - z)z,$ $x^2(x + u), x^2y, x^2z, (x - y)(x - z)u$	$(y, z), x + u$
57	$(x - y + z)z, (x - y)y, (x - u)u,$ $x^2(x + z), x^2y, x^2u, (x - y)(x - u)z$	$(y, u), x + z$
58	$z(x - y + z - u), y(x - y - u), u(x - u),$ $x^2(x + z), x^2y, x^2u, z(x - y)(x - u)$	$z, u, y + u$
61	$z(x - y + z), u(x - y + u), xy,$ $x(x + z)(x + u), y^3, z^3, u^3$	(y, z, u)
110	$(x - z + u)x, (y - z)z, (x - y)y,$ $u^3, u^2(y - z), u^2(x - y), xyz$	z, u

TABLE 28. Ideals and invariants when $(V(P), F(P)) = (8, 17)$

5.3.4. $(V(P), F(P)) = (8, 18)$. In this case, there are two smooth Fano polytopes as shown in Table 29. Their cohomology rings can be distinguished by c.v.e. over $\mathbb{Z}/3$ as shown in the following table.

ID	vertices of P from 5th	minimal nonfaces
53	$(-1, -1, 1, 1), (0, 1, -1, 0), (1, 0, 0, -1), (-1, -1, 0, 0)$	36, 47, 125, 128, 138, 156, 248, 257, 348, 567
55	$(-1, -1, 1, 1), (0, 1, -1, 0), (1, 0, 0, -1), (0, 0, -1, -1)$	36, 47, 125, 128, 138, 156, 248, 257, 348, 567

TABLE 29. $(V(P), F(P)) = (8, 18)$

ID	\mathcal{I}	c.v.e. $\mathbb{Z}/3$
53	$(x - y)y, (x - z)z, (x + u)^2x, u^3, (x - y)u^2,$ $(x + u)xy, (x - z)u^2, (x + u)xz, (x - y)(x - z)u, xyz$	u
55	$(x - y - u)y, (x - z - u)z, x^3, (x - y)(x - z)u, u^3,$ $x^2y, u^2(x - z), x^2z, u^2(x - y), xyz$	(x, u)

TABLE 30. Ideals and invariants when $(V(P), F(P)) = (8, 18)$

5.4. The case where $V(P) = 9$.

5.4.1. $(V(P), F(P)) = (9, 18)$. In this case, there are four smooth Fano 4-polytopes as shown in Table 31 and they are all combinatorially equivalent to a direct sum of a 2-simplex and a 6-gon. Using the data in Table 31, we obtain Table 32 which shows that the four cohomology rings are not isomorphic to each other.

ID	vertices of P from 5th	minimal nonfaces
27	$(-1, -1, 0, 2), (0, 0, -1, 1), (0, 0, 1, -1), (0, 0, -1, 0), (0, 0, 0, -1)$	36, 38, 39, 47, 48, 49, 67, 69, 78, 125
46	$(-1, -1, 1, 1), (0, 0, -1, 1), (0, 0, 1, -1), (0, 0, -1, 0), (0, 0, 0, -1)$	36, 38, 39, 47, 48, 49, 67, 69, 78, 125
119	$(-1, 0, 0, 1), (1, 0, 0, -1), (-1, 0, 0, 0), (0, -1, -1, 1), (0, 0, 0, -1)$	15, 17, 19, 46, 47, 49, 56, 59, 67, 238
122	$(-1, 0, 0, 1), (1, 0, 0, -1), (-1, 0, 0, 0), (0, 0, 0, -1), (0, -1, -1, 0)$	15, 17, 18, 46, 47, 48, 56, 58, 67, 239

TABLE 31. $(V(P), F(P)) = (9, 18)$

ID	\mathcal{I}	s.v.e. $\mathbb{Z}/2$	c.v.e. $\mathbb{Z}/3$
27	$y(y+u), u(y+u), v(z-u), z(2x-z-v),$ $u(2x+y-v), v(2x-z-v), yz, yv, zu, x^3$	$(y+u, z+v, u+v)$	$(y+u, x)$
46	$y(x-y-u), u(x-y-u), v(x+z-u), z(x-z-v),$ $u(x+y-v), v(x-z-v), yz, yv, zu, x^3$	$u+v$	x
119	$x(x+z), z(x+z), v(y-z), y(y-u+v),$ $z(x+u-v), v(y-u+v), xy, xv, yz, u^3$	$x+z$	$(x+z, u)$
122	$x(x+z), z(x+z), u(y-z), y(y+u),$ $z(x-u), u(y+u), xy, xu, yz, v^3$	$(x+z, y+u, z+u)$	all

TABLE 32. Ideals and invariants when $(V(P), F(P)) = (9, 18)$

5.4.2. $(V(P), F(P)) = (9, 20)$. In this case, there are 17 smooth Fano 4-polytopes as shown in Table 33 and they are all combinatorially equivalent to a direct sum of two 1-simplices and a 5-gon.

ID	vertices of P from 5th	minimal nonfaces
71	$(-1, 0, 0, 1), (0, -1, 0, 1), (0, 0, -1, 1), (0, 0, 1, -1), (0, 0, -1, 0)$	15, 26, 37, 39, 48, 49, 78
73	$(-1, 0, 0, 1), (0, -1, 0, 1), (0, 0, -1, 1), (0, 0, 1, -1), (0, 0, 0, -1)$	15, 26, 37, 39, 48, 49, 78
76	$(-1, 0, 0, 1), (0, -1, 0, 1), (0, 0, 1, -1), (0, 0, -1, 0), (0, 0, 0, -1)$	15, 26, 38, 39, 47, 49, 78
77	$(-1, 0, 0, 1), (0, -1, 0, 1), (0, 1, -1, 0), (0, 1, 0, -1), (0, -1, 0, 0)$	15, 26, 29, 37, 48, 49, 68
79	$(-1, 0, 0, 1), (0, -1, 0, 1), (0, 1, -1, 0), (1, 0, 0, -1), (-1, 0, 0, 0)$	15, 19, 26, 37, 48, 49, 58
81	$(-1, 0, 0, 1), (0, -1, 0, 1), (0, 1, -1, 0), (1, 0, 0, -1), (0, 0, 0, -1)$	15, 19, 26, 37, 48, 49, 58
82	$(-1, 0, 0, 1), (0, -1, 0, 1), (0, 1, -1, 0), (0, -1, 0, 0), (0, 0, 0, -1)$	15, 26, 28, 37, 48, 49, 69
84	$(-1, 0, 0, 1), (0, -1, 0, 1), (0, 1, 0, -1), (0, -1, 0, 0), (0, 0, -1, 0)$	15, 26, 28, 39, 47, 48, 67
88	$(-1, 0, 0, 1), (0, -1, 0, 1), (0, 1, 0, -1), (0, -1, 0, 0), (0, 1, -1, -1)$	15, 26, 28, 39, 47, 48, 67
90	$(-1, 0, 0, 1), (0, -1, 0, 1), (0, 1, 0, -1), (0, 0, -1, 0), (0, 0, 0, -1)$	15, 26, 29, 38, 47, 49, 67
91	$(-1, 0, 0, 1), (0, -1, 0, 1), (0, 1, 0, -1), (0, 0, 0, -1), (0, 1, -1, -1)$	15, 26, 28, 39, 47, 48, 67
92	$(-1, 0, 0, 1), (0, -1, 0, 1), (0, 1, 0, -1), (0, 0, 0, -1), (0, 0, -1, -1)$	15, 26, 28, 39, 47, 48, 67
102	$(-1, 0, 0, 1), (0, -1, 1, 0), (0, 1, -1, 0), (0, -1, 0, 0), (0, 0, 0, -1)$	15, 26, 28, 37, 38, 49, 67
103	$(-1, 0, 0, 1), (0, -1, 1, 0), (0, 1, -1, 0), (0, -1, 0, 0), (0, 1, -1, -1)$	15, 26, 28, 37, 38, 49, 67
107	$(-1, 0, 0, 1), (0, -1, 1, 0), (0, 0, -1, 0), (0, 0, 0, -1), (0, 0, -1, -1)$	15, 26, 37, 39, 48, 49, 78
113	$(-1, 0, 0, 1), (0, 0, 1, -1), (0, -1, 0, 0), (0, 0, -1, 0), (0, 0, 0, -1)$	15, 27, 38, 39, 46, 49, 68
120	$(-1, 0, 0, 1), (1, 0, 0, -1), (-1, 0, 0, 0), (0, -1, 0, 0), (0, 0, -1, 0)$	15, 17, 28, 39, 46, 47, 56

TABLE 33. $(V(P), F(P)) = (9, 20)$

We see that

$$\begin{aligned}
X_{73} &\cong X_{76} \cong X_{92}, & X_{77} &\cong X_{88}, & X_{81} &\cong X_{103}, \\
X_{82} &\cong X_{91} \cong X_{107}, & X_{90} &\cong X_{113}, & &
\end{aligned} \tag{5.7}$$

see Subsection 5.5. Using the data in Table 33, we obtain the following table.

ID	\mathcal{I}	s.v.e.	s.v.e. $\mathbb{Z}/2$	4-v.e. $\mathbb{Z}/2$
71	$x^2, y^2, z(z+v), v(z-u+v),$ $u(x+y-u), v(x+y-v), zu$	x, y		
73 (76, 92)	$x^2, y^2, z^2, v(z-u),$ $u(x+y-u-v), v(x+y-v), zu$	x, y, z		
77 (88)	$x^2, y(y-z+v), v(x+z-v),$ $z^2, u(x-u), v(x+y-u), yu$	$x, z, x-2u$	(x, z)	(x, z, u, v)
79	$x(x+v), v(v-y), y(y-z),$ $z^2, u(y-u), v(x+y-u), xu$	$z, z-2y$		
81 (103)	$x^2, v(x-u), y(y-z), z^2,$ $u(y-u-v), v(y-v), xu$	$x, z, z-2y$	(x, z)	(x, y, z, v)
82 (91, 107)	$x^2, y(y-z+u), u(y-z+u),$ $z^2, u(x+y-v), v(x-v), yv$	$x, z, x-2v,$ $z-2y-2u$	(x, z)	$(x, y+u, z, v)$
84	$x^2, y(y+u), u(x-u), v^2,$ $u(x+y-z), z(x-z), yz$	$x, v, x-2u,$ $x-2z$	(x, v)	all
90 (113)	$x^2, y^2, v(y-z), u^2,$ $v(x-v), z(x-z-v), yz$	$x, y, u,$ $x-2v$		
102	$x^2, y(y+u), u^2, z^2,$ $u(y-z), v(x-v), yz$	∞	(x, z, u)	
120	$x(x+z), z^2, u^2, v^2,$ $y^2, z(x-y), xy$	∞	(y, z, u, v)	

TABLE 34. Ideals and invariants when $(V(P), F(P)) = (9, 20)$

Table 34 shows that the ten cohomology rings in the table are distinguished by the invariants in the table except for 77 and 81. The s.v.e. over $\mathbb{Z}/2$ of $H^*(X_{77})$ and $H^*(X_{81})$ are both (x, z) . Therefore, if there is an isomorphism $H^*(X_{77}) \rightarrow H^*(X_{81})$, then it induces an isomorphism $(H^*(X_{77}) \otimes \mathbb{Z}/2)/(x, z) \rightarrow (H^*(X_{81}) \otimes \mathbb{Z}/2)/(x, z)$. However, Table 35 shows that this does not occur because the degree sequences of the ideals are different, so $H^*(X_{77}) \not\cong H^*(X_{81})$.

ID	$\mathcal{I} \otimes \mathbb{Z}/2 + (x, z)$
77(88)	$y(y+v), v^2, u^2, v(y+u), yu$
81(103)	$uv, y^2, u(y+u), v(y+v)$

TABLE 35. Distinguishment between $H^*(X_{77})$ and $H^*(X_{81})$

5.4.3. $(V(P), F(P)) = (9, 21)$. In this case, there are four smooth Fano 4-polytopes and there are two combinatorial types among them. Indeed, the combinatorial type of ID 52 is different from the others as is seen from Table 36.

ID	vertices of P from 5th	minimal nonfaces
51	$(-1, -1, 1, 1), (0, 1, -1, 0), (0, 1, 0, -1), (0, -1, 0, 0), (0, 0, -1, 0)$	28, 29, 36, 39, 47, 68, 125, 156, 157, 159, 348
52	$(-1, -1, 1, 1), (0, 1, -1, 0), (1, 0, 0, -1), (0, 0, -1, 0), (0, 0, 0, -1)$	19, 28, 36, 38, 47, 49, 125, 156, 257, 567
56	$(-1, -1, 1, 1), (0, 1, -1, 0), (0, -1, 0, 0), (0, 0, -1, 0), (0, 0, 0, -1)$	27, 28, 36, 38, 49, 67, 125, 156, 158, 159, 347
89	$(-1, 0, 0, 1), (0, -1, 0, 1), (0, 1, 0, -1), (0, -1, 0, 0), (1, 0, -1, -1)$	15, 26, 28, 47, 48, 67, 178, 239, 349, 359, 369

TABLE 36. $(V(P), F(P)) = (9, 21)$

Using the data in Table 36, we obtain Table 37. It shows that $H^*(X_{52})$ can be distinguished from the other three cohomology rings by the degree sequences of the ideals. On the other hand, Table 38 shows that the three cohomology rings are not isomorphic to each other.

ID	\mathcal{I}
51	$u(x-z+u), v(x-y-z+u), y(x-y-v), v(x-y-v),$ $z(x-z), yu, x^2(x+u), x^2y, x^2z, x^2v, u(x-v)(x-z)$
52	$v(x-z), u(x-y), y(x-y-u), u^2, z(x-z-v),$ $v^2, x^3, x^2y, x^2z, xyz$
56	$z(x+z), u(u+z), y(x-y-u), u(x-y-u), v(x-v),$ $yz, x^2(x+z), x^2y, x^2u, x^2v, z(x-u)(x-v)$
89	$x(x-v), y(y+u), u(y-z+u), z(x-z-v), u(x-u-v),$ $yz, zu(x-v), v^2(z-u), v^2(z+v), v^2x, v^2y$

TABLE 37. Ideals when $(V(P), F(P)) = (9, 21)$

ID	s.v.e. $\mathbb{Z}/2\mathbb{Z}$	c.v.e. $\mathbb{Z}/3\mathbb{Z}$	c.v.e. $\mathbb{Z}/2\mathbb{Z}$
51	\emptyset	$(z, y+v)$	$x+u, x+u+v, z, y+z, y+v, y+z+v$
52	(u, v)		
56	\emptyset	$(x+z, y+u, v)$	
89	\emptyset	$(x, y+u)$	$x, x+z+u, y+u, z+v, u+v$

TABLE 38. $(V(P), F(P)) = (9, 21)$

5.5. Diffeomorphism. In the previous subsections, we claimed some diffeomorphisms among toric Fano 4-folds, i.e. (5.1), (5.2), (5.3), (5.4), (5.5), (5.7). In this subsection, we establish them by showing that the condition in Lemma 2.3 is satisfied in each case.

In the following, we express the vertices of a smooth Fano polytope P_q with ID number q in terms of a matrix where each row shows a vertex of P_q and the numbers written on the left side of a matrix are the numbers of the vertices. For instance, the vertices of P_{70} are arranged as $(1, 2, 3, 4, 5, 6)$ while the vertices of P_{141} are arranged as $(1, 2, 3, 6, 5, 4)$ in their matrices. The correspondence $(1, 2, 3, 4, 5, 6) \rightarrow (1, 2, 3, 6, 5, 4)$ gives a bijection between the vertices of P_{70} and P_{141} , which preserves the combinatorial structures of P_{70} and P_{141} . The multiplication by a 4×4 unimodular matrix from the right shows a unimodular transformation. In each case, we will see that the resulting vectors of the corresponding vertices agree up to sign, so the condition in Lemma 2.3 is satisfied.

- 70 and 141

$$\begin{array}{c}
1 \\
2 \\
3 \\
4 \\
5 \\
6
\end{array}
\begin{array}{c}
\mathbf{70} \\
\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & -1 & 1 & 1 \\ 0 & 0 & -1 & -1 \end{pmatrix}
\end{array}
\begin{array}{c}
1 \\
2 \\
3 \\
6 \\
5 \\
4
\end{array}
\begin{array}{c}
\mathbf{141} \\
\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & -1 & -1 \\ -1 & -1 & 0 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}
\end{array}
\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \\ -1 & -1 & 1 & 1 \\ 0 & 0 & 1 & 1 \end{pmatrix}$$

- 30 and 43

$$\begin{array}{c}
1 \\
2 \\
3 \\
4 \\
5 \\
6 \\
7
\end{array}
\begin{array}{c}
\mathbf{30} \\
\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & -1 & 0 & 2 \\ 0 & 0 & -1 & 1 \\ 0 & 0 & 0 & -1 \end{pmatrix}
\end{array}
\begin{array}{c}
1 \\
2 \\
3 \\
4 \\
5 \\
7 \\
6
\end{array}
\begin{array}{c}
\mathbf{43} \\
\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & -1 & 0 & 2 \\ 0 & 0 & -1 & -1 \\ 0 & 0 & 0 & -1 \end{pmatrix}
\end{array}
\begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \\ 1 & 1 & 0 & -2 \\ 0 & 0 & -1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \\ 1 & 1 & 0 & -2 \\ 0 & 0 & -1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

- 68 and 134

$$\begin{array}{c}
1 \\
2 \\
3 \\
4 \\
5 \\
6 \\
7 \\
8 \\
9
\end{array}
\begin{array}{c}
\mathbf{77} \\
\begin{pmatrix}
1 & 0 & 0 & 0 \\
0 & 1 & 0 & 0 \\
0 & 0 & 1 & 0 \\
0 & 0 & 0 & 1 \\
-1 & 0 & 0 & 1 \\
0 & -1 & 0 & 1 \\
0 & 1 & -1 & 0 \\
0 & 1 & 0 & -1 \\
0 & -1 & 0 & 0
\end{pmatrix}
\end{array}
\begin{array}{c}
1 \\
6 \\
3 \\
4 \\
5 \\
2 \\
9 \\
8 \\
7
\end{array}
\begin{array}{c}
\mathbf{88} \\
\begin{pmatrix}
1 & 0 & 0 & 0 \\
0 & -1 & 0 & 1 \\
0 & 0 & 1 & 0 \\
0 & 0 & 0 & 1 \\
-1 & 0 & 0 & 1 \\
0 & 1 & 0 & 0 \\
0 & 1 & -1 & -1 \\
0 & -1 & 0 & 0 \\
0 & 1 & 0 & -1
\end{pmatrix}
\end{array}
\begin{pmatrix}
-1 & 0 & 0 & 0 \\
0 & 1 & 0 & -1 \\
0 & 0 & 1 & 0 \\
0 & 0 & 1 & 0 \\
0 & 0 & 0 & -1 \\
0 & 0 & 1 & 0 \\
0 & 0 & 0 & -1 \\
0 & 0 & 0 & -1 \\
0 & 0 & 0 & -1
\end{pmatrix}
=
\begin{pmatrix}
-1 & 0 & 0 & 0 \\
0 & -1 & 0 & 0 \\
0 & 0 & 1 & 0 \\
0 & 0 & 0 & -1 \\
1 & 0 & 0 & -1 \\
0 & 1 & 0 & -1 \\
0 & 1 & -1 & 0 \\
0 & -1 & 0 & 1 \\
0 & 1 & 0 & 0
\end{pmatrix}$$

• 81 and 103

$$\begin{array}{c}
1 \\
2 \\
3 \\
4 \\
5 \\
6 \\
7 \\
8 \\
9
\end{array}
\begin{array}{c}
\mathbf{81} \\
\begin{pmatrix}
1 & 0 & 0 & 0 \\
0 & 1 & 0 & 0 \\
0 & 0 & 1 & 0 \\
0 & 0 & 0 & 1 \\
-1 & 0 & 0 & 1 \\
0 & -1 & 0 & 1 \\
0 & 1 & -1 & 0 \\
1 & 0 & 0 & -1 \\
0 & 0 & 0 & -1
\end{pmatrix}
\end{array}
\begin{array}{c}
1 \\
6 \\
3 \\
4 \\
5 \\
2 \\
9 \\
8 \\
7
\end{array}
\begin{array}{c}
\mathbf{103} \\
\begin{pmatrix}
0 & 0 & 1 & 0 \\
0 & 0 & 0 & 1 \\
1 & 0 & 0 & 0 \\
0 & -1 & 1 & 0 \\
0 & -1 & 0 & 0 \\
0 & 1 & -1 & -1 \\
-1 & 0 & 0 & 1 \\
0 & 1 & 0 & 0 \\
0 & 1 & -1 & 0
\end{pmatrix}
\end{array}
\begin{pmatrix}
0 & 0 & 1 & 0 \\
-1 & 0 & 0 & 1 \\
-1 & 0 & 0 & 0 \\
0 & 1 & 0 & 0
\end{pmatrix}
=
\begin{pmatrix}
-1 & 0 & 0 & 0 \\
0 & 1 & 0 & 0 \\
0 & 0 & 1 & 0 \\
0 & 0 & 0 & -1 \\
1 & 0 & 0 & -1 \\
0 & -1 & 0 & 1 \\
0 & 1 & -1 & 0 \\
-1 & 0 & 0 & 1 \\
0 & 0 & 0 & 1
\end{pmatrix}$$

• 82 and 91 and 107

$$\begin{array}{c}
1 \\
2 \\
3 \\
4 \\
5 \\
6 \\
7 \\
8 \\
9
\end{array}
\begin{array}{c}
\mathbf{82} \\
\begin{pmatrix}
1 & 0 & 0 & 0 \\
0 & 1 & 0 & 0 \\
0 & 0 & 1 & 0 \\
0 & 0 & 0 & 1 \\
-1 & 0 & 0 & 1 \\
0 & -1 & 0 & 1 \\
0 & 1 & -1 & 0 \\
0 & -1 & 0 & 0 \\
0 & 0 & 0 & -1
\end{pmatrix}
\end{array}
\begin{array}{c}
1 \\
6 \\
3 \\
4 \\
5 \\
2 \\
9 \\
7 \\
8
\end{array}
\begin{array}{c}
\mathbf{91} \\
\begin{pmatrix}
1 & 0 & 0 & 0 \\
0 & -1 & 0 & 1 \\
0 & 0 & 1 & 0 \\
0 & 0 & 0 & 1 \\
-1 & 0 & 0 & 1 \\
0 & 1 & 0 & 0 \\
0 & 1 & -1 & -1 \\
0 & 1 & 0 & -1 \\
0 & 0 & 0 & -1
\end{pmatrix}
\end{array}
\begin{pmatrix}
-1 & 0 & 0 & 0 \\
0 & 1 & 0 & 0 \\
0 & 0 & 1 & 0 \\
0 & 0 & 1 & 0 \\
0 & 0 & 0 & -1 \\
0 & 0 & 0 & 0 \\
0 & 0 & 0 & -1 \\
0 & 0 & 0 & -1 \\
0 & 0 & 0 & 1
\end{pmatrix}
=
\begin{pmatrix}
-1 & 0 & 0 & 0 \\
0 & -1 & 0 & 0 \\
0 & 0 & 1 & 0 \\
0 & 0 & 0 & -1 \\
1 & 0 & 0 & -1 \\
0 & 1 & 0 & -1 \\
0 & 1 & -1 & 0 \\
0 & 1 & 0 & 0 \\
0 & 0 & 0 & 1
\end{pmatrix}$$

$$\begin{array}{c}
1 \\
3 \\
2 \\
8 \\
5 \\
9 \\
6 \\
7 \\
4
\end{array}
\begin{array}{c}
\mathbf{107} \\
\begin{pmatrix}
1 & 0 & 0 & 0 \\
0 & 0 & 1 & 0 \\
0 & 1 & 0 & 0 \\
0 & 0 & 0 & -1 \\
-1 & 0 & 0 & 1 \\
0 & 0 & -1 & -1 \\
0 & -1 & 1 & 0 \\
0 & 0 & -1 & 0 \\
0 & 0 & 0 & 1
\end{pmatrix}
\end{array}
\begin{pmatrix}
-1 & 0 & 0 & 0 \\
0 & 0 & -1 & 0 \\
0 & -1 & 0 & 0 \\
0 & 0 & 0 & 1
\end{pmatrix}
=
\begin{pmatrix}
1 & 0 & 0 & 0 \\
0 & -1 & 0 & 0 \\
0 & 0 & -1 & 0 \\
0 & 0 & 0 & -1 \\
-1 & 0 & 0 & 1 \\
0 & 1 & 0 & -1 \\
0 & -1 & 1 & 0 \\
0 & 1 & 0 & 0 \\
0 & 0 & 0 & 1
\end{pmatrix}$$

• 90 and 113

$$\begin{array}{c}
1 \\
2 \\
3 \\
4 \\
5 \\
6 \\
7 \\
8 \\
9
\end{array}
\begin{array}{c}
\mathbf{90} \\
\begin{pmatrix}
1 & 0 & 0 & 0 \\
0 & 1 & 0 & 0 \\
0 & 0 & 1 & 0 \\
0 & 0 & 0 & 1 \\
-1 & 0 & 0 & 1 \\
0 & -1 & 0 & 1 \\
0 & 1 & 0 & -1 \\
0 & 0 & -1 & 0 \\
0 & 0 & 0 & -1
\end{pmatrix}
\end{array}
\begin{array}{c}
1 \\
6 \\
2 \\
9 \\
5 \\
8 \\
3 \\
7 \\
4
\end{array}
\begin{array}{c}
\mathbf{113} \\
\begin{pmatrix}
1 & 0 & 0 & 0 \\
0 & 0 & 1 & -1 \\
0 & 1 & 0 & 0 \\
0 & 0 & 0 & -1 \\
-1 & 0 & 0 & 1 \\
0 & 0 & -1 & 0 \\
0 & 0 & 1 & 0 \\
0 & -1 & 0 & 0 \\
0 & 0 & 0 & 1
\end{pmatrix}
\end{array}
\begin{pmatrix}
1 & 0 & 0 & 0 \\
0 & 0 & 1 & 0 \\
0 & -1 & 0 & 1 \\
0 & 0 & 0 & 1
\end{pmatrix}
=
\begin{pmatrix}
1 & 0 & 0 & 0 \\
0 & -1 & 0 & 0 \\
0 & 0 & 1 & 0 \\
0 & 0 & 0 & -1 \\
-1 & 0 & 0 & 1 \\
0 & 1 & 0 & -1 \\
0 & -1 & 0 & 1 \\
0 & 0 & -1 & 0 \\
0 & 0 & 0 & 1
\end{pmatrix}$$

6. c_1 -PRESERVING COHOMOLOGY RING ISOMORPHISM

As we observed, cohomology rings do not distinguish toric Fano manifolds as varieties. Very recently, motivated by McDuff's question on the uniqueness of toric actions on a monotone symplectic manifold, Y. Cho, E. Lee, S. Park and the third author made the following conjecture and verified it for Fano Bott manifolds ([7]).

Conjecture ([7]). *If there is a cohomology ring isomorphism between toric Fano manifolds which preserves their first Chern classes, then they are isomorphic as varieties.*

In this section, we prove the following theorem mentioned in Introduction, which gives further supporting evidence to the conjecture.

Theorem 6.1. *The conjecture above is true for toric Fano d -folds with $d = 3, 4$ or with Picard number $\geq 2d - 2$.*

In order to prove the theorem above, it suffices to check that there is no c_1 -preserving cohomology ring isomorphism between toric Fano d -folds which have isomorphic cohomology rings. If there is a c_1 -preserving cohomology ring isomorphism between toric Fano d -folds X and Y , then $c_1(X)^d$ evaluated on the fundamental class of X , in other words the degree $(-K_X)^d$ of X , agrees with that of Y . We obtain the following tables from the database of Øbro. They together with Tables 1 and 6 show that the degrees are different for toric Fano 3- or 4-folds which have isomorphic cohomology rings except one pair ID 70 and 141.

ID	11, 18	10, 13
degree	52, 44	44, 40

TABLE 39. Degrees of toric Fano 3-folds with isomorphic cohomology rings

ID	70, 141	30, 43	68, 134	129, 136	28, 32	67, 118	123, 125
degree	513, 513	592, 400	432, 480	496, 400	478, 382	351, 447	415, 367

ID	74, 96	83, 108	95, 131	29, 39	111, 116	50, 57	73, 76, 92
degree	480, 352	448, 352	416, 352	463, 337	389, 347	417, 369	394, 330, 310

ID	77, 88	81, 103	82, 91, 107	90, 113	72, 87	78, 86
degree	405, 331	373, 325	341, 363, 229	352, 320	308, 268	298, 278

TABLE 40. Degrees of toric Fano 4-folds with isomorphic cohomology rings

As for ID 70 and 141, more detailed observation is necessary. It follows from Table 9 and Remark 2.2 that

$$\begin{aligned} H^*(X_{70}) &= \mathbb{Z}[x, y]/(x^3, y(x-y)^2), & c_1(X_{70}) &= x + 3y \\ H^*(X_{141}) &= \mathbb{Z}[x, y]/(x^3, y^2(x-y)), & c_1(X_{141}) &= 2x + 3y. \end{aligned} \quad (6.1)$$

An elementary computation shows that an isomorphism $H^*(X_{70}) \rightarrow H^*(X_{141})$ is given by either $(x, y) \rightarrow (x, x-y)$ or $(x, y) \rightarrow (-x, -x+y)$ but both isomorphisms are not c_1 -preserving. This completes the proof of the theorem when $d = 3, 4$.

As investigated in Section 3, toric Fano d -folds with Picard number $\geq 2d - 2$ which have isomorphic cohomology rings are the following three pairs:

- (1) $Z_3^d = Z_3 \times (X_{P_6})^{\frac{d-3}{2}}$ and $Z_4^d = Z_4 \times (X_{P_6})^{\frac{d-3}{2}}$, where d is odd ≥ 3
- (2) $W_5^d = W_5 \times (X_{P_6})^{\frac{d-4}{2}}$ and $W_7^d = W_7 \times (X_{P_6})^{\frac{d-4}{2}}$, where d is even ≥ 4 .
- (3) $W_6^d = W_6 \times (X_{P_6})^{\frac{d-4}{2}}$ and $W_8^d = W_8 \times (X_{P_6})^{\frac{d-4}{2}}$, where d is even ≥ 4 ,

see (3.6) and (3.7). Here

$$(Z_3, Z_4) = (X_{10}, X_{13}), \quad (W_5, W_7) = (X_{72}, X_{87}), \quad (W_6, W_8) = (X_{78}, X_{86})$$

as mentioned above Convention in Sections 4 and 5. The degree of a product of projective varieties X and Y of dimension p and q is $\binom{p+q}{p}$ times the product of degrees of X and Y , so it follows from Tables 39 and 40 that the three pairs above have different degrees respectively. This completes the proof of the theorem.

REFERENCES

- [1] B. Assarf, M. Joswig and A. Paffenholz, Smooth Fano polytopes with many vertices, *Discrete Comput. Geom.* **52** (2014), 153–194.
- [2] V. V. Batyrev, On the classification of smooth projective toric varieties, *Tohoku Math J.* **43** (1991), 569–585.
- [3] V. V. Batyrev, On the classification of toric Fano 4-folds, *J. Math. Sci. (New York)* **94** (1999), 1021–1050.
- [4] V. Buchstaber and T. Panov, Toric Topology, Mathematical Surveys and Monographs, **204**. American Mathematical Society, Providence, RI, 2015.
- [5] V. M. Buchstaber, N. Yu. Erokhovets, M. Masuda, T. E. Panov, and S. Park, Cohomological rigidity of manifolds defined by 3-dimensional polytopes, *Uspekhi Mat. Nauk* **72** (2017), no. 2(434), 3–66.
- [6] C. Casagrande, The number of vertices of a Fano polytope, *Ann. Inst. Fourier* **56** (2006), 121–130.
- [7] Y. Cho, E. Lee, M. Masuda, and S. Park, Unique toric structure on a Fano Bott manifold, arXiv:2005.02740.
- [8] S. Choi, Classification of Bott manifolds up to dimension 8, *Proc. Edinb. Math. Soc. (2)* **58** (2015), no. 3, 653–659.
- [9] S. Choi, M. Masuda and S. Murai, Invariance of Pontrjagin classes for Bott manifolds, *Alg. & Geom. Top.* **15** (2015) 965–986.
- [10] S. Choi, M. Masuda and D. Y. Suh, Quasitoric manifolds over a product of simplices, *Osaka J. Math.* **47** (2010), no. 1, 109–129.
- [11] S. Choi, M. Masuda and D. Y. Suh, Topological classification of generalized Bott towers, *Trans. Amer. Math. Soc.* **362** (2010), no. 2, 1097–1112.
- [12] S. Choi, M. Masuda and D. Y. Suh, Rigidity problems in toric topology: a survey, *Proc. Steklov Inst. Math.* **275** (2011), no. 1, 177–190.
- [13] M. W. Davis, When are two Coxeter orbifolds diffeomorphic?, *Mich. Math. J.* **63** (2014), 401–421.
- [14] M. W. Davis and T. Januszkiewicz, Convex polytopes, Coxeter orbifolds and torus actions, *Duke Math. J.* **62** (1991), 417–451.
- [15] W. Fulton, “Introduction to Toric Varieties”, Ann. of Math. Studies 131, Princeton Univ. Press, 1993.
- [16] S. Hasui, H. Kuwata, M. Masuda, and S. Park, Classification of toric manifolds over an n -cube with one vertex cut, *IMRN*, ny161, <https://doi.org/10.1093/imrn/rny161>, Published: 05 July 2018. (Online Published) arXiv:1705.07530.
- [17] M. Masuda, Equivariant cohomology distinguishes toric manifolds, *Adv. Math.* **218** (2008), 2005–2012.
- [18] M. Masuda and T. Panov, Semi-free circle actions, Bott towers, and quasitoric manifolds, *Mat. Sb.* **199** (2008), no. 8, 95–122.
- [19] M. Masuda and D. Y. Suh, Classification problems of toric manifolds via topology, *Contemp. Math.*, **460**, Amer. Math. Soc., Providence, RI, 2008.
- [20] M. Øbro, An algorithm for the classification of smooth Fano polytopes, arXiv:0704.0049v1.
- [21] M. Øbro, Classification of terminal simplicial reflexive d -polytopes with $3d - 1$ vertices. *Manuscr. Math.* **125**(1), 69–79 (2008).
- [22] T. Oda, “Convex Bodies and Algebraic Geometry -An introduction to the theory of toric varieties”, *Ergeb. Math. Grenzgeb. (3)*, Vol. 15, Springer-Verlag, Berlin, Heidelberg, New York, London, Paris, Tokyo, 1988.
- [23] H. Sato, Towards the classification of higher-dimensional toric Fano varieties, *Tohoku Math J.* **52** (2000), 383–413.
- [24] B. Sturmfels, “Gröbner Bases and Convex Polytopes,” volume 8 of University Lecture Series, American Mathematical Society, Providence, RI, 1995.
- [25] K. Watanabe and M. Watanabe, The classification of Fano 3-folds with torus embeddings, *Tokyo J. Math.* **5** (1982), no. 1, 37–48.

(A. Higashitani) DEPARTMENT OF PURE AND APPLIED MATHEMATICS, GRADUATE SCHOOL OF INFORMATION SCIENCE AND TECHNOLOGY, OSAKA UNIVERSITY, SUITA, OSAKA 565-0871, JAPAN
E-mail address: `higashitani@ist.osaka-u.ac.jp`

(K. Kurimoto) DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, KYOTO SANGYO UNIVERSITY, KYOTO 603-8555, JAPAN
E-mail address: `i1885045@cc.kyoto-su.ac.jp`

(M. Masuda) DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, OSAKA CITY UNIVERSITY, SUGIMOTO, SUMIYOSHI-KU, OSAKA 558-8585, JAPAN
E-mail address: `masuda@sci.osaka-cu.ac.jp`